

Alphabetical Index

Alnus sinuata , 4
 Alnus tenuifolia , 44
 Alnus viridis crispa , 15
 Asclepias incarnata , 10
 Athyrium niponicum , 15

 Betula pubescens , 4

 Caltha leptosepala , 13
 Caltha leptosepala howellii , 14
 Caltha palustris , 14
 Convallaria majalis , 6
 Cornus sericea , 6
 Crataegus acclivis , 20
 Crataegus altaica , 26
 Crataegus anomala , 30
 Crataegus aprica , 51
 Crataegus armoldiana , 45
 Crataegus azarolus , 47
 Crataegus canbyi , 35
 Crataegus champlainensis , 48
 Crataegus chrysocharpa , 29
 Crataegus coccinoides , 29
 Crataegus columbiana , 30
 Crataegus crus , 33
 Crataegus cuneata , 49
 Crataegus douglasii , 19
 Crataegus durobrivensis , 22
 Crataegus ellwangeriana , 46
 Crataegus flabellata , 27
 Crataegus flava , 50
 Crataegus heterophylla , 54
 Crataegus holmesiana , 19
 Crataegus hupehensis , 52
 Crataegus intricata , 31
 Crataegus jackii , 36
 Crataegus jonesiae , 35
 Crataegus laciniata , 48
 Crataegus laevigata , 25
 Crataegus maximowiczii , 32
 Crataegus missouriensis , 16
 Crataegus mollis , 20
 Crataegus monogyna , 28
 Crataegus nigra , 54
 Crataegus opaca , 23
 Crataegus pedicellata , 16
 Crataegus pedicellata gloriosa , 24
 Crataegus pennsylvanica , 45
 Crataegus pentagyna , 51
 Crataegus phaenopyrum , 33
 Crataegus pinnatifida , 50
 Crataegus pinnatifida major , 47
 Crataegus pruinosa , 27
 Crataegus punctata , 23
 Crataegus rivularis , 36
 Crataegus rotundifolia , 25
 Crataegus sanguinea , 34
 Crataegus schraderana , 46
 Crataegus submollis , 21
 Crataegus succulenta , 22
 Crataegus uniflora , 37
 Crataegus x grignonensis , 13

 Eriophorum angustifolium , 38

 Filipendula kamtschatica , 13
 Filipendula multijuga , 57
 Filipendula ulmaria , 11

 Hosta longissima , 37

 Iris sibirica , 43

 Juncus effusus , 43

 Leucopum aestivum , 43

Lysimachia nummularia , 44
 Lysimachia vulgaris , 42
 Lythrum salicaria , 13

 Malus coronaria , 24
 Malus florentina , 34
 Mentha aquatica , 52
 Mimulus guttatus , 55
 Mimulus lewisii , 38

 Picea glauca , 5
 Picea glehnii , 32
 Populus tremula , 8
 Primula denticulata , 38

 Quercus bicolor , 18
 Quercus petraea , 53
 Quercus robur , 17

 Rosa canina , 10

 Salix acutifolia , 40
 Salix aegyptiaca , 56
 Salix alaxensis , 15
 Salix alba , 7
 Salix arenaria , 57
 Salix brachycarpa , 42
 Salix caprea , 39
 Salix commutata , 42
 Salix daphnoides , 40
 Salix fragilis , 39
 Salix hookeriana , 56
 Salix pentandra , 8
 Salix purpurea , 41
 Salix viminalis , 41
 Salix x mollissima hippophaefolia , 57

 Thuja occidentalis , 12
 Thuja plicata , 55

 Viburnum opulus , 9

Plants For a Future 2021CE Hardiness Zone 5, Wet Soil, Heavy Clay, Edibility

>= 1

pfaf.org

2021

Contents

1 Introduction	3
1.1 Pronunciation guide	3
1.2 vocabulary	3
2 2 tlin	4
2.1 bwon	4
2.2 mran	4
3 3 tlin	9
4 4 tlin	15
4.1 bwon	15
4.2 mran	16
5 5 tlin	44
5.1 bwon	44
5.2 mran	45
6 Alphabetical Index	57

1 Introduction

This is a book of plant information harvested from pfaf.org the goal is for future generations to have access to this valuable plant information. This book does Not contain any plant identification material, so you will need to use other references for actually identifying the plant. This book does contain information on how to grow the plants and their edible and other uses.

The plants are sorted by nitrogen-fixing ability, edibility, plant height, other uses, medicinal uses, and latin name. They start with nitrogen fixers because those are typically the easiest to establish pioneer plants.

Instructions for food forest design are in separate books. But basic idea is to have thorny plants on the border as a form of fence to keep out predators and animals that want to raid your food forest, with the shortest plants closest to the border, then gradually getting taller plants as you go in. For example starting with thorny herbaceous, then row of thorny shrub, the row of thorny trees. Afterwards can put in several rows of your staple crop, which typically is a variety of nut, acorn and bean producing tall trees. Then can put in a long rotation coppice of hardwood lumber trees you find useful, then a row or more of fruit trees, then a short rotation coppice to use for firewood, then some rows of berry, bean and nut shrubs, with the closest circle near the community being vegetables and perennials that take more care than the others.

You need about a hectare of land (2.5 acres) per person if you have 6 months growing season (where average temperature is above 10C), at least half of which should be dedicated to food, the other half can be to lumber, but can also be food trees that are occasionally used as lumber. A general formula is 6/monthsGrowingSeason to see how many hectares you need. This assumes you have at least 600mm of rainfall per year, and only works if you have more than 3 months growing season, as very little you can grow in less than that.

If you live in a dry area, especially one prone to flash floods, then it is imperative that you spend your dry season digging swales and vernal pools, that will collect the rain water during the flash floods and keep it in place so that it enters the ground, then you can grow things downhill of your vernal pools and swales. It is generally more important that they be extensive than deep. You want to ideally catch every drop, so that none of it leaves your property.

By contrast if you live in a very wet area with more than enough water, then you want to have ditches to help the water leave the property and not waterlog your soil.

In a medium wetness area you can use either method to create various microclimates for different kinds of plants.

In either case you want to start with about 1 in three plants being a nitrogen fixer (bwon), as it will help your food forest grow faster.

Summaries are given in the Cosmic Language (KsomPyash). Long descriptions are in English.

1.1 Pronunciation guide

a /a/, b /b/, c /ʃ/ "sh", d /d/, e /e/, f /f/, g /g/, h /h/, i /i/, j /ʒ/ "zh", k /k/, l /l/, m /m/, n /n/, o /o/, p /p/, q /ŋ/ "ng", r /r/, s /s/, t /t/, u /u/, v /v/, w /w/, x /x/, y /j/, z /z/, 2 /l/ (low tone), 6 /ə/ "uh", 7 /l/ (high tone).

1.2 vocabulary

tlin /tlin/ cold hardiness, cyuc /tju/ heat hardiness, kcak /kʌk/ food rating, tcan /tʃan/ other uses rating, dwat /dwat/ medicinal rating, hkuc /hku/ adult height, hsi2n /hsi:n/ adult width.

pcim /pʃim/ full (shade), hmim /hmim/ medium (shade), htut /htut/ no (shade), cyut /tʃut/ shade. li /li/ realis-mood (end of sentence). ta /ta/ topic-case.

hgat /hɡat/ drought (tolerant), kyuc /kju/ dry, tcup /tʃup/ damp, hja7n /hʒa:n/ wet, hs6s /hs6s/ puddle, hwic /hwif/ water (requirement).

hyis /hjis/ poor (soil), hras /hras/ sandy, jlut /ʒlut/ loamy, nyik /njik/ mud (silt/clay loam), frip /frip/ pottery (heavy clay), hpa7t /hpa7t/ topsoil.

frus /frus/ corrosive (very acid), flat /flat/ acid, prik /prik/ passive (neutral), lwis /lwis/ alkaline, slip /slip/ slippery (very alkaline), hsom /hsom/ salt (saline), hpa7tmwat /hpa7tmwat/ topsoil taste.

fyif /fjif/ self (fertile), xre7t /xre7t/, nyanhjen /njanhʒen/ male-female (dioecious, distinct male and female plants), hjon /hʒon/ sex.

hmuç /hmuç/ purpose, pcis /pʃis/ danger, trot /trot/ temperature, prop /prop/ fertile, tcan /tʃan/ other, hyum /hjum/ uses, dwat /dwat/ medicine, drut /drut/ description, kcac /kʌc/ region, tfo2n /tfo:n/ nitrogen-fixing, xlutkfam /xlut,kfam/ plant quantity, bwon /bwon/ bean, mran /mran/ corn/grain non-nitrogen-fixing plant, nrak /nrak/ continent (of origin), ksac /ksaç/ circumstances/habitat.

5 hmuctrotkcec
 hja7n hmuchwic
 frip hmuchpa7t
 102 xlutkfam

2 2 tlin

2.1 bwon

2.1.1 Alnus sinuata

Sitka Alder
tfo2n 2 tlin 9 cyuc 1 kcaak 3 tcan 1 dwat 4 m hkuc
hmimhtut cyutli tucphja7n hwicli hyishrlutnyikfripr hpa7tli flat-
prikrlwis hpa7tmwatli hjonli

drutta Alnus sinuata is a deciduous Shrub growing to 4 m (13ft 1in) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 2. It is in flower from April to June, and the seeds ripen from July to August. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. It can fix Nitrogen. Suitable for: medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista The freshly harvested inner bark is emetic but is alright once it has been dried[172].

kcakta Edible Parts: FlowersEdible Uses: Catkins - raw or cooked. A bitter flavour[172].

tcanta Fuel Pioneer Soil stabilization Tannin WoodThis is an excellent pioneer species for re-establishing woodlands on disused farmland, difficult sites etc[226]. Its fast rate of growth means that it quickly provides sheltered conditions to allow more permanent woodland trees to become established. In addition, bacteria on the roots fix atmospheric nitrogen - whilst this enables the tree to grow well in quite poor soils it also makes some of this nitrogen available to other plants growing nearby. Alder trees also have a heavy leaf canopy and when the leaves fall in the autumn they help to build up the humus content of the soil. Alder seedlings do not compete well in shady woodland conditions and so this species gradually dies out as the other trees become established[K]. The bark and the strobils are a source of tannin[82]. The tree has an extensive root system and can be planted to control banks from erosion[226, 229]. Wood - soft, straight-grained, very durable in water[82]. The tree is too small to be of use as a source of lumber, but the wood is used locally for fuel[226, 229].

dwatta Astringent Emetic Haemostatic Stomachic TonicThe bark is astringent, emetic, haemostatic, stomachic and tonic[172].

nrakta Western N. America - Alaska to California - east to Idaho and Montana. ksacta Moist woods, stream banks, margins of ponds, to open but moist montane slopes at or near the timber line[60].

propta Seed - best sown in a cold frame as soon as it is ripe and only just covered[200]. Spring sown seed should also germinate successfully so long as it is not covered[200, K]. The seed should germinate in the spring as the weather warms up. When large enough to handle, prick the seedlings out into individual pots. If growth is sufficient, it is possible to plant them out into their permanent positions in the summer, otherwise keep them in pots outdoors and plant them out in the spring. If you have sufficient quantity of seed, it can be sown thinly in an outdoor seed bed in the spring[78]. The seedlings can either be planted out into their permanent positions in the autumn/winter, or they can be allowed to grow on in the seed bed for a further season before planting them. Cuttings of mature wood, taken as soon as the leaves fall in autumn, outdoors in sandy soil.

nwatta Prefers a heavy soil and a damp situation[1, 11]. Grows well in heavy clay soils[11]. Tolerates very infertile sites[200]. This species is closely related to *A. viridis* and is included as a sub-species of *A. viridis* by many botanists[11]. A fast-growing but short-lived species, rarely living longer than 50 years[229]. It is a pioneer species of logged or burnt land, often forming dense thickets in low wet areas[229]. This species has a symbiotic relationship with certain soil micro-organisms, these form nodules on the roots of the plants and fix atmospheric nitrogen. Some of this nitrogen is utilized by the growing plant but some can also be used by other plants growing nearby[200]. In garden design, as well as the above-ground architecture of a plant, root structure considerations help in choosing plants that work together for their optimal soil requirements including nutrients and

water. The root pattern is branching: a heart root, dividing from the crown into several primary roots going down and out [2-1].

2.2 mran

2.2.1 Betula pubescens

White Birch, Downy birch

2 tlin 9 cyuc 3 kcaak 5 tcan 3 dwat 20 m hkuc
htut cyutli kyuctucphja7n hwicli hyishrasjlutnyikfripr hpa7tli frusflatprikrlwis hpa7tmwatli hjonli

drutta Betula pubescens is a deciduous Tree growing to 20 m (65ft) by 10 m (32ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 1. It is in flower in April, and the seeds ripen from August to September. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very acid soils. It cannot grow in the shade. It prefers dry moist or wet soil. The plant can tolerate strong winds but not maritime exposure.

pcista The aromatic and aliphatic hydrocarbons in birch tar are irritating to the skin. Do not use in patients with oedema or with poor kidney or heart functions [301]

kcakta Edible Parts: Flowers Inner bark Leaves SapEdible Uses: Tealner bark - cooked or dried, ground into a powder then used with cereals for making bread etc[2, 15, 46]. Inner bark is generally only seen as a famine food, used when other forms of starch are not available or are in short supply[177, K]. Sap - raw or cooked. A sweet flavour[2, 15, 177]. Harvested in early spring, before the leaves unfurl, by tapping the trunk. The flow is best on sunny days following a heavy frost. The sap is often concentrated into a sugar by boiling off the water. Between 4 and 7 litres can be drawn off a mature tree in a day and this will not kill the tree so long as the tap hole is filled up afterwards[115]. However, prolonged or heavy tapping will kill the tree. A beer can be fermented from the sap. An old English recipe for the beer is as follows: "To every Gallon of Birch-water put a quart of Honey, well stirr'd together; then boil it almost an hour with a few Cloves, and a little Limon-peel, keeping it well scum'd. When it is sufficiently boil'd, and become cold, add to it three or four Spoonfuls of good Ale to make it work...and when the Test begins to settle, bottle it up. . . it is gentle, and very harmless in operation within the body, and exceedingly sharpens the Appetite, being drunk ante pastum."[269]. Young leaves - raw or cooked[15, 177]. Young catkins[15]. No more details are given. A tea is made from the leaves[15] and another tea is made from the essential oil in the inner bark[21].

tcanta Adhesive Besom Charcoal Compost Dye Essential Fibre Fungicide Miscellaneous Paper Pioneer Polish Repellent Tannin Thatching Waterproofing WoodThe bark is used to make drinking vessels, canoe skins, roofing tiles etc. It is waterproof, durable, tough and resinous[11, 61]. Only the outer bark is removed, this does not kill the tree. It is most easily removed in late spring to early summer. The bark was pressed flat and stored until the following spring. When required for making canoes it would be heated over a fire to make it pliable for shaping to the canoe frame[257]. A pioneer species, it readily invades old fields, cleared or burnt-over land and creates conditions suitable for other woodland trees to become established. Since it is relatively short-lived and intolerant of shade, it is eventually out-competed by these trees[11, 186]. A tar-oil is obtained from the white bark in spring. It has fungicidal properties and is also used as an insect repellent[4, 14, 61, 100]. It makes a good shoe polish[61]. Another report says that an essential oil is obtained from the bark and this, called 'Russian Leather' has been used as a perfume[245]. A glue is made from the sap. Cordage can be made from the fibres of the inner bark. This inner bark can also be separated into thin layers and used as a substitute for oiled paper[4]. A decoction of the inner bark is used to preserve cordage, it is rich in tannin. The bark contains up to 16% tannin[223]. A brown dye is obtained from the inner bark. An oil similar to Wintergreen oil (obtained from *Gaultheria procumbens*) is obtained from the

produced in catkins early in the year; they are pollinated by bees and other insects rather than by the wind[11]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

5.2.23 Filipendula multijuga

5 tlin 9 cyuc 1 kcaak 0 tcan 0 dwat m hkuc
hmimhtut cyutli tucphja7n hwicli hrasjlutnyikfripr hpa7tli prikrlwis hpa7tmwatli fyifxre7t hjonli

drutta Filipendula multijuga is a PERENNIAL growing to 1.2 m (4ft). See above for USDA hardiness. It is hardy to UK zone 6. It is in flower from July to August. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies, beetles. The plant is self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcakta Edible Parts: LeavesEdible Uses: Young shoot tips - cooked[116].

nrakta E. Asia - Japan. ksacta Mountains, C. and S. Japan[58].

propta Seed - best sown in the autumn in a cold frame[1]. The seed can also be sown in a cold frame in spring, germinating best at a temperature of 10 - 13°C[200]. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer if they have grown enough. If not, keep them in a cold frame for the winter and plant them out in late spring. Division in autumn or winter[200]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring.

nwatta Requires a humus-rich moist soil in semi-shade[200]. Succeeds in full sun only if the soil is reliably moist throughout the growing season[200]. Dislikes dry or acid soils. Does well in marshy soils. Grows well in heavy clay soils. A very ornamental plant[116]. The flowers are very attractive to bees. Members of this genus are rarely if ever troubled by browsing deer[233].

5.2.24 Salix arenaria

5 tlin 9 cyuc 1 kcaak 3 tcan 2 dwat 1 m hkuc
htut cyutli tucphja7n hwicli hrasjlutnyikfripr hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta Salix arenaria is a deciduous Shrub growing to 1 m (3ft 3in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate maritime exposure.

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - raw or cooked. They are not very palatable[172].

tcanta Soil stabilizationThe plant has a spreading root system and is an effective soil binder on sand dunes[200]. Dynamic accumulator.

dwatta Anodyne FebrifugeThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

6 Alphabetical Index

nrakta W. Europe. ksacta Not known

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

5.2.25 Salix x mollisima hippophaeifolia

5 tlin 9 cyuc 1 kcaak 3 tcan 2 dwat m hkuc
htut cyutli tucphja7n hwicli hrasjlutnyikfripr hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta Salix x mollisima hippophaeifolia is a deciduous Shrub growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil.

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - cooked. Not very palatable[172].

tcanta BasketryThe stems are very flexible and are used in basket making[131]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. Succeeds in wet, ill-drained or intermittently flooded soils[1, 11]. Dynamic accumulator.

dwatta Anodyne FebrifugeThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta A hybrid of garden origin, *S. triandra* x *S. viminalis*. ksacta Not known in a truly wild situation.

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. This is a hybrid species and it will not breed true from seed. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Plants in this genus are notably susceptible to honey fungus[200]. Plants are rarely bisexual[200], but they are usually dioecious and then male and female plants must be grown if seed is required.

been used as a counter-irritant for the skin[257]. A poultice of the inner bark has been applied to carbuncles[257]. The bark has been pounded until it is as soft as cotton and then used to rub the face[257]. The very soft bark has been used to bind wounds and cover dressings[257]. The shredded bark has been used to cauterize sores and swellings[257].

nrakta Western N. America - British Columbia to California. ksacta Grows singly and in small groves in moist to swampy soils, usually in shady forests, from sea-level to 1350 metres[60, 82, 99].

propta Seed - best sown when ripe in the autumn in a cold frame[113]. Stored seed germinates best if given a short cold stratification[113]. It can be sown in a cold frame in late winter. When they are large enough to handle, prick the seedlings out into individual pots and grow them on in the greenhouse for their first winter. Plant them out into their permanent positions in late spring or early summer, after the last expected frosts. If growing large quantities of plants, the seed can be sown in an outdoor seed bed in mid spring[78]. Grow the plants on for two years and then plant them out into their permanent positions in late autumn or early spring. Cuttings of half-ripe wood, 5 - 8cm with a heel, July/August in a shaded frame. Forms roots by the end of September but it should be overwintered in a frame[78]. Cuttings of almost ripe wood, 5 - 10cm with a heel, September in a cold frame. Forms roots in the following summer. Plant out in autumn or spring[78].

nwatta Landscape Uses:Pest tolerant, Screen, Specimen. Prefers a moist loamy soil[1, 81]. Dislikes dry soils[11]. Not suitable for light sandy soils because it is shallow rooting[75], but it succeeds on most other soil types, even on waterlogged soils and heavy clays[81, 200]. Succeeds in chalky soils[200]. Prefers some shelter from the wind according to some reports[1, 81] whilst others say that it is fairly wind resistant[11, 75]. Trees tolerate salt winds[75] but not severe maritime exposure[K]. The wood and foliage are highly aromatic[226]. The foliage has a strong fruity smell, like pear-drops or crushed apples[185]. A fast growing tree, it often puts out new growth of 1 metre a year and can average 60cm a year for the first 40 years. Increase in girth can also be rapid, 5cm a year for the first 60 years has been recorded[185]. Sometimes planted for forestry in Britain[17], it is at its best in the moister western half of the country[11, 185]. There are many named varieties selected for their ornamental value[188]. Tolerant of light trimming, so long as this is not back into old wood, plants also retain their lower branches unless these are shaded out by neighbouring plants[11]. The foliage turns bronze in cold weather[81]. Trees are notably susceptible to honey fungus[81, 200]. Special Features: North American native, Inconspicuous flowers or blooms.

5.2.21 Salix aegyptiaca

5 tin 9 cyuc 1 kcak 2 tcan 2 dwat 4 m hkuc htut cyutli tcuphja7n hwicli hrasljutnyikfrp hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta Salix aegyptiaca is a deciduous Shrub growing to 4 m (13ft) by 5 m (16ft). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower from January to February. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required). and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil.

kcakta Edible Parts: Flowers Inner bark LeavesEdible Uses: Drinkinrner bark - raw or cooked. It can be dried, ground into a powder and added to cereal flours for use in making bread etc. A famine food, it is only used when all else fails[172]. Young shoots - cooked. They are not very palatable[172]. Male catkins - sugared[11, 177]. A perfumed drink is made from the catkins[11].

tcanta Dynamic accumulator.

dwatta Anodyne FebrifugeThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta N. Africa to W. Asia. ksacta Mountains in Asia[50].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Closely related to *S. cinerea*[1]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

5.2.22 Salix hookeriana

Dune Willow

5 tin 9 cyuc 1 kcak 3 tcan 2 dwat 1 m hkuc htut cyutli tcuphja7n hwicli hrasljutnyikfrp hpa7tli flatprikhsom hpa7tmwatli nyanhjen hjonli

drutta Salix hookeriana is a deciduous Shrub growing to 1 m (3ft 3in) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower from April to May, and the seeds ripen in June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required). and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils and can grow in saline soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate maritime exposure.

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - raw or cooked. They are not very palatable[172]. The leaves have been used as a flavouring in cooked foods[257].

tcanta Basketry Fibre Hair Miscellany WoodStems are very flexible and are used in basket making[61, 118]. Tolerates maritime exposure[11, 200]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. The bark has been used extensively in basket making[257]. Fibres from the inner bark can be twisted into long ropes[226]. The soft roots have been used as a towel to rub down after bathing[257]. An infusion of the roots has been used as a hair wash[257]. Wood - light, soft, close grained[82]. Dynamic accumulator.

dwatta Anodyne Antidote Febrifuge MiscellanyThe leaves have been used as an antidote to shellfish poisoning[257]. The fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta Western N. America - Alaska to California. ksacta Borders of salt marshes and ponds, also on sandy coastal dunes[82]. Streams, ponds and sloughs near the shore[229].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Tolerates maritime exposure[11, 200]. A fast-growing but short-lived species[229]. Hybridizes freely with other members of this genus[200]. Although the flowers are

inner bark[21, 61]. It is used medicinally and also makes a refreshing tea[21]. The young branches are very flexible and are used to make whisks, besoms etc[6]. They are also used in thatching and to make wattles[4]. The leaves are a good addition to the compost heap, improving fermentation[20]. A black paint is obtained from the soot of the plant[61]. A high quality charcoal is obtained from the bark. It is used by artists, painters etc. Wood - soft, light, durable. It is used for a wide range of purposes including furniture, tool handles, carving, toys etc[100, 238]. It is a source of charcoal that is used by artists and is also pulped and used for making paper[238].

dwatta Antirheumatic Antiseborrheic Astringent Bitter Diaphoretic Diuretic Eczema Lithotropic Miscellany Skin UrinaryAnti-inflammatory, cholagogue, diaphoretic[21, 165, 201]. The bark is diuretic and laxative[7]. The inner bark is bitter and astringent, it is used in treating intermittent fevers[4]. An oil obtained from the inner bark is astringent and is used in the treatment of various skin afflictions, especially eczema and psoriasis[4, 238]. The bark is usually obtained from trees that have been felled for timber and can be distilled at any time of the year[238]. The buds are balsamic[7]. The young shoots and leaves secrete a resinous substance which has acid properties, when combined with alkalis it is a tonic laxative[4]. The leaves are anticholesterolemic and diuretic[7]. They also contain phytosides, which are effective germicides[7]. An infusion of the leaves is used in the treatment of gout, dropsy and rheumatism, and is recommended as a reliable solvent of kidney stones[4]. The young leaves and leaf buds are harvested in the spring and dried for later use[238]. A decoction of the leaves and bark is used for bathing skin eruptions[4]. The vernal sap is diuretic[4]. The boiled and powdered wood has been applied to chafed skin[257]. Moxa is made from the yellow fungous excrescences of the wood, which sometimes swell out of the fissures[4]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve *Betula* species for infections of the urinary tract, kidney and bladder stones, rheumatism (see [302] for critics of commission E).

nrakta Most of Europe, including Britain, east to W. Siberia and central Asia. ksacta Open woodland and heaths, usually on acid soils, from sea level to 830 metres[1, 17, 100].

propta Seed - best sown as soon as it is ripe in a light position in a cold frame[78, 80, 113, 134]. Only just cover the seed and place the pot in a sunny position[78, 80, 134]. Spring sown seed should be surface sown in a sunny position in a cold frame[113, 134]. If the germination is poor, raising the temperature by covering the seed with glass can help[134]. When they are large enough to handle, prick the seedlings out into individual pots and grow them on in a cold frame for at least their first winter. Plant them out into their permanent positions in late spring or early summer, after the last expected frosts. If you have sufficient seed, it can be sown in an outdoor seedbed, either as soon as it is ripe or in the early spring - do not cover the spring sown seed. Grow the plants on in the seedbed for 2 years before planting them out into their permanent positions in the winter[78, 80, 113, 134].

nwatta Succeeds in a well-drained light loamy soil in a sunny position[11, 200]. Grows well in heavy clay soils. Tolerates a wet position[11, 24], succeeding in poorly drained soils[186]. Fairly wind tolerant[200]. Prefers an acid soil. A very ornamental tree and fast growing, capable of growing 1 metre a year but it is short-lived[186]. It is one of the first trees to colonize open land and it creates a suitable environment for other woodland trees to follow[11]. These trees eventually shade out the birch trees[186]. Trees take about 15 years from seed to produce their own seed[98]. Although closely related, it does not usually hybridize with *B. pendula*[11]. It hybridizes freely with *B. pendula* according to another report[186]. A superb tree for encouraging wildlife, it has over 200 associated insect species[24, 30]. A good plant to grow near the compost heap, aiding the fermentation process[14, 20]. It is also a good companion plant, its root activity working to improve the soil[14]. Trees are notably susceptible to honey fungus[200].

2.2.2 Picea glauca

White Spruce, Black Hills Spruce, Canadian Spruce 2 tin 6 cyuc 2 kcak 3 tcan 2 dwat 15 m hkuc

htut cyutli tcuphja7n hwicli hyishrasljutnyikfrp hpa7tli frusflatprik hpa7tmwatli hjonli

drutta *Picea glauca* is an evergreen Tree growing to 15 m (49ft) by 5 m (16ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 2. It is in leaf all year, and the seeds ripen in September. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid and neutral soils and can grow in very acid soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure. It cannot tolerate atmospheric pollution.

kcakta Edible Parts: Flowers Inner bark Seed Seed-podEdible Uses: Condiment Gum TeaYoung male catkins - raw or cooked. Used as a flavouring[172]. Immature female cones - cooked. The central portion, when roasted, is sweet and syrupy[172]. The cones are about 5cm long[82]. Inner bark - raw or cooked[257]. It can be dried, ground into a powder and then used as a thickener in soups etc or added to cereals when making bread[172]. Usually harvested in the spring[257], it is an emergency food that is only used when all else fails. Seed - raw[172]. The seed is about 2 - 4mm long[229] and is too small and fiddly to be worthwhile unless you are desperate[172]. A refreshing tea, rich in vitamin C, can be made from the young shoot tips[172]. The trunk yields a gum, used for chewing[183, 257]. Spruce oil, distilled from the leaves and twigs, is used in the food industry to flavour chewing gum, ice cream, soft drinks and sweets[183].

tcanta Baby care Dye Gum Musical Pitch Repellent Shelterbelt Striving Tannin Waterproofing WoodA fairly wind resistant tree, it can be grown as part of a shelterbelt planting[200]. The cultivar 'Denstat' has been recommended[200]. The leaves have been burnt to repel insects[257]. Various native North American Indian tribes made a string from the long roots of this species and used it to stitch the bark of their canoes and to make baskets etc[226, 257]. The rotten, dried, finely powdered wood has been used as a baby powder and as a treatment for skin rashes[257]. The bark is a source of tannin[226]. A yellow-brown dye can be obtained from the rotten wood[257]. The pitch obtained from the trunk can be used as a waterproofing sealant in canoes[257]. Wood - straight-grained, resilient, light, soft, not strong. Used for construction and as a source of pulp for paper making[46, 61, 171, 229]. The resonance of the wood, and its capacity to transmit vibrations, make it an ideal wood for guitars, violins, piano soundboards etc[226].

dwatta Antiseptic Digestive Diuretic Laxative Pectoral Poultice Salve Skin TB Urinary Vitamin CWhite spruce was widely employed medicinally by several native North American Indian tribes, who valued it especially for treating chest complaints[257]. It is little, if at all, used in modern herbalism. An infusion of the cones has been used in the treatment of urinary troubles[257]. The inner bark is pectoral[257]. It has been chewed, and an infusion drunk, in the treatment of TB, influenza, coughs and colds[257]. An infusion is also drunk in the treatment of rheumatism[257]. The inner bark has also been used as a poultice on sores and infected areas, and has also been used to bandage cuts[257]. The tea made from the young shoot tips has antiseptic properties[226]. It is used in the treatment of respiratory infections[257]. A decoction of the stems is used as a herbal steam bath in the treatment of rheumatism[257]. The gum is antiseptic, digestive, laxative, pectoral and salve[257]. A decoction has been used in the treatment of respiratory complaints[257]. The gum obtained from the trunk (probably pitch[K]) has been used as a salve on sores and cuts[257]. A poultice of the gum mixed with oil has been used to treat skin rashes, scabies, persistent scabs, growing boils etc, and has also been used on wounds where there is blood poisoning[257]. The rotten, dried, finely powdered wood has been used as a baby powder and as a treatment for skin rashes[257].

nrakta Northern N. America - Alaska to Newfoundland. ksacta Woods in good soils[43]. Along streams and lakes and also on rocky hills and slopes, succeeding in a variety of soil conditions[229].

propta Seed - stratification will probably improve germination so sow fresh seed in the autumn in a cold frame if possible[80]. Sow stored seed as early in the year as possible in a cold frame[78]. A position in light shade is probably best[78]. Seed should not be allowed to dry out and should

be stored in a cool place[80]. Prick out the seedlings into individual pots when they are large enough to handle and grow them on in the greenhouse or cold frame for their first winter. They can be planted out into their permanent positions in early summer of the following year, or be placed in an outdoor nursery bed for a year or so to increase in size. They might need protection from spring frosts. Cuttings of semi-ripe terminal shoots, 5- 8cm long, August in a frame. Protect from frost. Forms roots in the spring[78]. Cuttings of mature terminal shoots, 5 - 10cm long, September/October in a cold frame. Takes 12 months[78]. Cuttings of soft to semi-ripe wood, early summer in a frame. Slow but sure.

nwatta Landscape Uses:Christmas tree, Firewood, Screen, Specimen. Likes abundant moisture at the roots, if grown in drier areas it must be given a deep moist soil[11]. Tolerates poor peaty soils[200]. Succeeds in wet cold and shallow soils but is not very wind-firm in shallow soils[1]. Prefers a pH between 4 to 6[200]. Dislikes shade[200]. Intolerant of atmospheric pollution[11]. Resists wind exposure to some degree[200]. A fast growing tree, especially when young with annual increases of up to 1 metre in height[185]. New growth takes place from April to July. Growth slows considerably as the trees grow older[185]. It is an important forestry tree in N. America and is also planted for timber in N. Europe. It is sometimes used as a 'Christmas tree', but is unsited for this because its leaves quickly fall[226]. Seed production begins at approximately 20 years, though reliable crops make take twice that long[229]. Heavy crops are produced every 2 - 5 years[229]. Trees should be planted into their permanent positions when they are quite small, between 30 and 90cm. Larger trees will check badly and hardly put on any growth for several years. This also badly affects root development and wind resistance[200]. In some upland areas, especially over granitic or other base-poor soils, growth rate and health have been seriously affected by aluminium poisoning induced by acid rain[200]. Plants are strongly outbreeding, self-fertilized seed usually grows poorly[200]. They hybridize freely with other members of this genus[200]. There are some named varieties selected for their ornamental value[200]. The crushed leaves are quite aromatic. Some people find the smell distasteful saying that it is like skunk[226], whilst others say it has a pleasant smell like blackcurrants or mouldy grapefruit[185]. Special Features:North American native, Inconspicuous flowers or blooms.

2.2.3 Cornus sericea

Red Osier Dogwood, Western dogwood

2 tlin 7 cyuc 2 kcaak 4 tcan 2 dwat 2.5 m hkuc hmimhtut cyutli tcuphja7n hwicli hrasjlutnyikfrip hpa7tli flat-prikwlwis hpa7tmwatli xre7t hjonli

drutta Cornus sericea is a deciduous Shrub growing to 2.5 m (8ft) by 4 m (13ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 2 and is not frost tender. It is in flower from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcakta Edible Parts: Fruit Oil SeedEdible Uses: Oil OilFruit - raw or cooked[105, 161, 257]. Juicy[101]. Bitter and unpalatable according to some reports[2], it was mixed with other fruits such as juneberries (Amelanchier spp) and then dried for winter use by native North Americans[257]. The fruit can cause nausea[172]. The fruit is up to 9mm in diameter[200]. Seed[101]. No more details are given, but the seeds are quite small and woody, looking rather less than edible[K]. An edible oil is obtained from the seed[4].

tcanta Basketry Dye Fibre Fodder Miscellany Oil Oil TeethA fibre obtained from the bark is used as cordage[99]. The bark can be twisted into a rope[257]. The powdered bark has been used as a toothpowder to preserve the gums and keep the teeth white[4]. An oil obtained from the seed burns well and can be used in lighting[4]. A red dye can be obtained from the bark mixed with cedar ashes[257]. The branches are pliable, they are used as rims in basket making[99, 257]. The stem wood is very tough and flexible[212]. Plants can be grown as a tall ground cover for colonising large areas.

The cultivar 'Flaviramea' has been recommended[208]. Red osier is noted as a good Carbon Farming Solution plant. It can be coppiced, used as fodder and act as a living fence. It is a industrial biomass crop.

dwatta Analgesic Astringent Refrigue Miscellany Poul-tice Purgative Skin Stimulant TonicRed osier dogwood was widely employed by several native North American Indian tribes who valued it especially for its astringent and tonic bark, using it both internally and externally to treat diarrhoea, fevers, skin problems etc[257]. It is little used in modern herbalism. The bark and the root bark are anal-gesic, astringent, febrifuge, purgative, slightly stimulant and tonic[4, 172, 257]. Drying the bark removes its tendency to purge[172]. A decoction has been used in the treatment of headaches, diarrhoea, coughs, colds and fevers[257]. Externally, the decoction has been used as a wash for sore eyes, styes and other infections and also to treat skin complaints such as poison ivy rash and ulcers[257]. The bark shavings have been applied as a dressing on wounds to stop the bleed-ing[257]. A poultice of the soaked inner bark, combined with ashes, has been used to alleviate pain[257]. The plant is said to have cured hydrophobia[4].

nrakta N. America - Newfoundland to New York, west to Alaska and California. Naturalized in Britain[17]. ksacta Shores and thickets[43]. Along streams, rivers and moist sites, 450 - 2700 metres[229].

propta Seed - best sown as soon as it is ripe in a cold frame or in an outdoors seedbed if there is sufficient seed[80, 113]. The seed must be separated from the fruit flesh since this contains germination inhibitors[80, 164]. Stored seed should be cold stratified for 3 - 4 months and sown as early as possible in the year[164]. Scarification may also help as may a period of warm stratification before the cold stratifi-cation[80, 164]. Germination, especially of stored seed, can be very slow, taking 18 months or more[164]. Prick out the seedlings of cold-frame sown seeds into individual pots as soon as they are large enough to handle and grow the plants on for their first winter in a greenhouse, planting out in the spring after the last expected frosts. Cuttings of half-ripe side shoots, July/August in a frame. Cuttings of mature wood of the current year's growth, taken with a heel if possible, autumn in a cold frame. High percentage[78]. Layering of new growth in June/July. Takes 9 months[78].

nwatta Agroforestry Services: Living fence Fodder: Bank Industrial Crop: Biomass Management: Cop-pice Regional CropAn easily grown plant, it succeeds in any soil of good or moderate fertility[1], ranging from acid to shallow chalk[200]. Grows well in heavy clay soils. Prefers a moist soil and a position in sun or partial shade[108]. Succeeds in poorly drained soils[200]. Plants are hardy to about -35°C[184]. A rampant suckering shrub[1]. A number of cul-tivars have been developed for their ornamental value[182]. This species is closely allied to *C. alba*[11]. The flowers are very attractive to bees[108]. Plants in this genus are notably resistant to honey fungus[200]. The plant is heat tolerant in zones 8 through 1. (Plant Hardiness Zones show how well plants withstand cold winter temperatures. Plant Heat Zones show when plants would start suffering from the heat. The Plant Heat Zone map is based on the number of "heat days" experienced in a given area where the temperature climbs to over 86 degrees F (30°C). At this temperature, many plants begin to suffer physiological damage. Heat Zones range from 1 (no heat days) to 12 (210 or more heat days). For example Heat Zone. 11-1 indicates that the plant is heat tolerant in zones 11 through 1. For polyculture design as well as the above-ground architecture (form - tree, shrub etc. and size shown above) information on the habit and root pattern is also useful and given here if available. The plant growth habit is multistemmed with multiple stems from the crown [1-2].

2.2.4 Convallaria majalis

Lily Of The Valley, European lily of the valley

2 tlin 7 cyuc 1 kcaak 3 tcan 3 dwat m hkuc pcmhmmimhtut cyutli kyuctcuphja7n hwicli hyishrasjlutnyik-frip hpa7tli flatprikwlwis hpa7tmwatli fyifxre7t hjonli

drutta Convallaria majalis is a PERENNIAL growing to 0.2 m (0ft 8in) by 0.3 m (1ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 3 and is not frost tender. It is in flower from May to June, and the seeds ripen

5.2.19 Mimulus guttatus

Yellow Monkey Flower, Seep monkeyflower

5 tlin 9 cyuc 2 kcaak 0 tcan 2 dwat 0.6 m hkuc hmimhtut cyutli tcuphja7nhs6s hwicli hrasjlutnyikfrip hpa7tli flatprikwlwis hpa7tmwatli hjonli

drutta Mimulus guttatus is a PERENNIAL growing to 0.6 m (2ft) by 0.6 m (2ft in). See above for USDA hardiness. It is hardy to UK zone 6. It is in flower from July to September. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can grow in water.

kcakta Edible Parts: LeavesEdible Uses: Leaves - raw or cooked[94, 105, 155, 172]. A slightly bitter flavour, they are added to salads[161, 183]. The leaves are used as a lettuce substitute[257].

dwatta Astringent Bach Poultice VulneryA The plant is astringent, poultice and vulnery[94, 172, 257]. A decoction of the leaves and stems has been used as a herbal steam bath for chest and back soreness[257]. A poultice of the crushed leaves has been applied to wounds, rope burns etc[257]. The plant is used in Bach flower remedies - the keywords for pre-scribing it are 'Fear or anxiety of a known origin'[209].

nrakta Western N. America - Alaska to Mexico. Natural-ized in Britain. ksacta Streams and wet places below 3000 metres[187]. Banks of streams in Britain, where it appears quite native[17].

propta Seed - sow spring in a greenhouse and only lightly cover the seed[200]. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer. If you have sufficient seed it can also be sown in situ in late spring after the last ex-pected frosts[200]. Division in spring[200]. Very easy, large divisions can be planted out direct into their permanent po-sitions. We have found that it is better to pot up the smaller divisions and grow them on in light shade in a cold frame un-til they are well established before planting them out in late spring or early summer. Cuttings of softwood, 5cm long in spring or summer[200].

nwatta Prefers rather moist places such as damp bor-ders, margins of streams and boggy spots[1, 111]. Grows well in heavy clay soils. Thrives in full sun so long as the soil does not dry out[200], otherwise it is best grown in partial shade[187]. Grows well in shallow water, but it can be inva-sive[200]. Hardy to at least -20°C[187]. There is some confu-sion over the naming of this plant, the entry on edibility may refer to *M. luteus*, which is closely related to this species.

5.2.20 Thuja plicata

Western Red Cedar, Giant Arborvitae, Giant Cedar, Incense Cedar, Western Red Cedar

5 tlin 8 cyuc 1 kcaak 0 tcan 2 dwat 60 m hkuc hmimhtut cyutli tcuphja7n hwicli hrasjlutnyikfrip hpa7tli flat-prikwlwisslip hpa7tmwatli hjonli

drutta Thuja plicata is an evergreen Tree growing to 60 m (196ft) by 12 m (39ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 6. It is in leaf all year, in flower from March to April, and the seeds ripen from Septem-ber to October. The species is monoecious (individual flow-ers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and ba-sic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil. The plant can tolerate maritime exposure.

kcakta Edible Parts: Inner barkEdible Uses: GumInner bark - fresh or dried[2, 105, 161]. The inner bark can be dried, ground into a powder and then used as a thickening in soups etc or mixed with cereals when making bread. A pitch obtained from the trunk has been used as a chewing gum[257].

tcanta Baby Care Basketry Brush Dye Fibre Fuel Fungi-cide Gum Hair Hedge Insulation Paper Roofing Scourer Shelterbelt Stuffing Thatching Tinder Wick Wood-Tolerant of light trimming and of reasonable exposure, this

species can be grown as a hedge or as part of a shelter-belt[75]. An infusion of the boughs can be used as a hair wash to treat dandruff and scalp germs[257]. The fibrous inner bark can be pounded until it is soft and then used as a sponge for scouring dishes etc, or can be used for making rough clothing, blankets, mats, ropes, sanitary towels, a padding in a baby's cradle, nappies etc. Waterproof hats, capes, trousers, skirts etc can be made from the inner bark[257]. It is also used in thatching and as a stuffing material for mat-tresses[46, 61, 82, 99, 118, 171, 226, 257]. Inner bark strips have been used as a roofing material[257]. The bark has also been used to make paint brushes[257]. The inner bark has been used to make a wick for oil lamps[257]. The inner bark has been used for making baskets[257]. The roots are used in basket making, making nets etc[99]. The roots have been used in coiled and imbricated baskets[257]. The roots have been peeled, split and used to make coiled watertight bas-kets that can be used for boiling water[257]. The roots are harvested in the spring or early autumn when it is easier to remove the bark. The outer strips of the roots are used to make the bottom of the basket, the centre core is used in the coils and the root bark, because of its toughness, is used to make the edges[257]. The fibrous bark is used for roofing and the sides of shelters. It is also used as an insulation[61, 99]. A fibre obtained from the bark is used in making paper. The fibre is about 3.8mm long (this refers to the heartwood fibre, the inner bark fibre is probably longer)[189]. Branches can be harvested at any time of the year, they are cut into us-able pieces and pre-soaked in clear water prior to cooking. They are then cooked for six hours or more with lye. It is difficult to rinse it to clear water because it seems to be a dye material[189]. The fibre is then hand pounded with mal-lets, or put through a blender or a ball mill for six hours. It is difficult to hydrate properly. The resulting paper is a rich deep brown/red[189]. The slender pliable branches are used as a high quality rope[257]. They are gathered in spring, peeled and, if thick, are split into halves or quarters. They are then twisted and worked until soft and pliable and finally woven together to make the rope[99]. A green dye can be obtained from the leaves and twigs[257]. The inner bark can be used as a tinder[257]. Wood - aromatic, light, soft, straight-grained, not strong, very resistant to decay. This resistance to decay is probably due to the existence of powerful fungi-cides in the wood[226]. The wood from fallen trees remains sound for at least 100 years[226]. It is pale to dark red in colour[226]. The wood was widely utilized by many native North American Indian tribes who used it for making a wide range of items including canoes, houses, totem poles, bows, spoons, ladders and tools[226, 257]. It is currently used in making greenhouses[1, 11, 82, 171]. The wood is not of such good quality when grown in mild humid areas[1]. It makes a good fuel, burning with very little smoke, though it burns quickly[99].

dwatta Abortifacient Analgesic Antidandruff An-tirheumatic Astringent Emmenagogue Febrifuge Kidney Odontalgic Pectoral Poultice Skin Stomachic TB VDWestern red cedar was employed medicinally by a number of native North American Indian tribes, who used it to treat a wide range of complaints[257]. It is seldom, if ever, used in mod-ern herbalism. An infusion of the leaves has been used in the treatment of stomach pains and diarrhoea[257]. A decoction of the leaves has been used in the treatment of colds[257]. A decoction of the powdered leaves has been used externally to treat various internal pains, including rheumatism[257]. The leaf buds have been chewed in the treatment of toothaches and sore lungs[257]. A decoction of the buds has been used as a gargle[257]. A decoction of the small branches has been used in the treatment of coughs, colds and tuberculosis[257]. A weak infusion has been drunk in the treatment of painful joints caused by rheumatism or arthritis[257]. A poultice of the crushed bough tips and oil has been applied to the back and chest in the treatment of bronchitis, rheumatism, stom-ach pains and swollen neck[257]. An infusion of the twigs has been used as a wash in the treatment of venereal disease sores[257]. A decoction of the boughs has been used as an antidandruff shampoo[257]. A decoction of the stem tips and the roots has been used in the treatment of colds[257]. An infusion of the bark and twigs has been used in the treat-ment of kidney complaints[257]. An infusion of the seeds and twigs has been used in the treatment of fevers[257]. The chewed bark, or a decoction of the bark, has been drunk to induce menstruation[257]. A moxa of the inner bark has

5.2.17 Crataegus heterophylla

5 tlin 9 cyuc 2 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykrip hpa7tli flatprikwis hpa7tmwatli xre7t hjonli

drutta Crataegus heterophylla is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardness. It is hardy to UK zone 6 and is not frost tender. It is in flower from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[177]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is up to 1.5cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Europe to E. Asia - Spain to China. Rarely self-sown in Britain. ksacta Found in secondary georgian oak forests.

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A very ornamental plant[1].

5.2.18 Crataegus nigra

Hungarian Hawthorn

5 tlin 9 cyuc 2 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykrip hpa7tli flatprikwis hpa7tmwatli xre7t hjonli

drutta Crataegus nigra is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardness. It is hardy to UK zone 6 and is not frost tender. It is in flower in May. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[177]. A soft flesh[200]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is about 1cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta E. and C. Europe - Hungary. ksacta Woods[50].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies. The plant is self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in full shade (deep woodland) semi-shade (light woodland) or no shade. It prefers dry moist or wet soil.

pcista All parts of the plant are poisonous[7, 10, 19]. However, the toxic principle is very poorly absorbed when taken orally so poisoning is unlikely to occur[65]. The leaves can be a mild skin irritant[22]. Overdose may lead to nausea, vomiting, stupor, colour perception disorders, and cardiac arrhythmias. Internal use preparations no longer considered safe [301].

kcakta A wine can be prepared from the flowers, mixed with raisins[183].

tcanta Dye EssentialAn essential oil is obtained from the flowers[46, 171]. It is used in perfumery and for snuff[238]. A green dye is obtained from the leaves in spring[13, 14, 115]. A yellow dye is obtained from the leaves in autumn[14]. Plants can be grown as a ground cover in woodland shade or in a shrubbery[200, 208]. As a garden ornamental it has gained the Royal Horticultural Society's Award of Garden Merit.

dwatta Antianxiety Antiarrhythmic Anticonvulsant Antispasmodic Cardiotonic Diuretic Emetic Febrifuge Laxative Poulitice SedativeLily of the valley has a long and proven reputation in herbal medicine in the treatment of heart complaints. It contains the glycosides convallarin and convallamarin which are powerful cardiac tonics and diuretics and are also used in allopathic medicine[244]. However, because of the plants potential toxic properties it should never be used without expert advice[9]. All parts of the plant are antispasmodic, cardiotonic, strongly diuretic, emetic, febrifuge, laxative and sedative[4, 7, 9, 21, 46, 165, 222, 254]. The plant is usually harvested when in flower and can be dried for later use[4], though it is stronger acting when fresh[238]. The inflorescence is said to be the most active medicinally and is often harvested separately[4]. An infusion of the flowers and roots is a digitalis substitute (obtained from Digitalis species), though less powerful, that is especially useful in the treatment of valvula heart diseases, cardiac debility, dropsy and chronic lung problems such as emphysema[4, 222, 254]. Lily of the valley encourages the heart to beat more slowly, regularly and efficiently, at the same time it is strongly diuretic, reducing blood volume and lowering blood pressure[254]. Its effect is less cumulative than digitalis which makes it safer for elderly patients[238]. It is often prescribed combined with the fruits of Crataegus spp[238]. An ointment made from the roots is used in the treatment of burns and to prevent scar tissue[222]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve Convallaria majalis : Lily Of The Valley for arrhythmia, cardiac insufficiency, nervous heart complaints (see [302] for critics of commission E).

nrakta Europe, including Britain, from Scandinavia to Spain east to N.E. Asia, Greece and the Caucasus. ksacta Dry shady woodland, usually on calcareous soils[7, 13, 17], and especially in ash woodlands[4].

propta Seed - best sown as soon as it is ripe, otherwise in late winter, in a cold frame[164]. Germination, particularly of stored seed can be very slow, taking 2 - 12 months or more at 15°C[164]. Sow the seed thinly so that the seedlings can be allowed to grow on undisturbed in the pot for their first year. Apply a liquid feed during the growing season to ensure that the seedlings are well fed. Divide the young plants into individual pots when they die down in late summer and grow them on in pots in a shady position in a cold frame for at least another year before planting them out into their permanent positions when they are dormant[K]. Division in September[111]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring.

nwatta Landscape Uses:Border, Container, Erosion control, Ground cover, Massing, Specimen, Woodland. garden Succeeds in almost any situation, including the dense dry shade of large trees[1, 4, 24]. Prefers a position in semi-shade in a moderately fertile well-drained moist woodland soil[200, 208]. Grows well in heavy clay, sand or chalky soils[208]. Dislikes pure clay soils and boggy sites[233].

Plants are hardy to -20°C or lower[200]. A polymorphic species[200]. It is a very ornamental plant, though it can become very invasive once it is established[28]. Plants can take a couple of years to become established[208]. There are several named varieties, selected for their ornamental value[233]. The flowers are sweetly scented[245]. Lily of the valley is occasionally cultivated as a medicinal plant for herbalists and allpaths. Plants seem to be immune to the predations of rabbits[233]. A good bee plant[4]. Special Features: Not North American native, Naturalizing, All or parts of this plant are poisonous, Suitable for cut flowers, Fragrant flowers.

2.2.5 Salix alba

White Willow

2 tlin 8 cyuc 1 kcaak 4 tcan 3 dwat 25 m hkuc htut cyutli tucphja7n hwicli hrasjlutnykrip hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta Salix alba is a deciduous Tree growing to 25 m (82ft) by 10 m (32ft) at a fast rate. See above for USDA hardness. It is hardy to UK zone 2 and is not frost tender. It is in flower from April to May, and the seeds ripen in June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate maritime exposure.

pcista Gastrointestinal bleeding and kidney damage possible. Avoid concurrent administration with other aspirin-like drugs. Avoid during pregnancy. Drug interactions associated with salicylates applicable [301].

kcakta Edible Parts: Inner bark LeavesEdible Uses: TeaInner bark - raw or cooked. It can be dried, ground into a powder and added to cereal flour then used in making bread etc[2]. A very bitter flavour, especially when fresh[2, 115], it is used as a famine food when all else fails[172]. Leaves and young shoots - raw or cooked[2, 177]. Not very palatable[172]. They are used only in times of scarcity[105]. The leaves can be used as a tea substitute[61].

tcanta Basketry Charcoal Paper Pioneer Shelterbelt String WoodLandscape Uses: Aggressive surface roots possible, Pollard, Specimen. The young stems are very flexible and are used in basket making[13, 46, 61]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. The bark can be used for tying plants[61]. A fibre obtained from the stems is used in making paper[189]. The stems are harvested in spring or summer, the leaves are removed and the stems steamed until the fibres can be stripped. The fibres are cooked for 2 hours with lye and then beaten with mallets or put through a blender. The paper is red/brown in colour[189]. A fast growing tree and tolerant of maritime exposure, it can be grown as a shelterbelt[75]. The plant's rapid growth and wind tolerance make it a very good pioneer species to use in establishing woodland conditions in difficult sites. Spacing cuttings about every 5 metres will soon provide shelter and a suitable environment for planting out woodland trees that are not so wind tolerant. The main disadvantage in using this species is that the roots are far-ranging and the plant is quite greedy, so it will not as much effect as species such as the alders (Alnus species) in enriching the soil and thus feeding the woodland plants[K]. Wood - elastic, soft, easy to split, does not splinter. Used for construction, turnery, poles, tool handles etc[11, 46, 61]. The wood is also used to make charcoal[11], which has medicinal uses[7]. Dynamic accumulator.

dwatta Anodyne Antiinflammatory Antiperiodic Antirheumatic Antiseptic Astringent Diaphoretic Diuretic Dysentery Febrifuge Hypnotic Sedative TonicJustly famous as the original source of salicylic acid (the precursor of aspirin), white willow and several closely related species have been used for thousands of years to relieve joint pain and manage fevers[254]. The bark is anodyne, anti-inflammatory, antiperiodic, antiseptic, astringent, diaphoretic, diuretic, febrifuge, hypnotic, sedative and tonic[4, 7, 9, 21, 165]. It has been used internally in the treatment of dyspepsia

connected with debility of the digestive organs[4], rheumatism, arthritis, gout, inflammatory stages of auto-immune diseases, feverish illnesses, neuralgia and headache[238]. Its tonic and astringent properties render it useful in convalescence from acute diseases, in treating worms, chronic dysentery and diarrhoea[4]. The fresh bark is very bitter and astringent[222]. It contains salicin, which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[213]. The bark is harvested in the spring or early autumn from 3 - 6 year old branches and is dried for later use[7, 9]. The leaves are used internally in the treatment of minor feverish illnesses and colic[238]. An infusion of the leaves has a calming effect and is helpful in the treatment of nervous insomnia[7]. When added to the bath water, the infusion is of real benefit in relieving widespread rheumatism[7]. The leaves can be harvested throughout the growing season and are used fresh or dried[238]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve *Salix* / Willow for diseases accompanied by fever, rheumatic ailments, headaches (see [302] for critics of commission E).

nrakta Europe, including Britain, from Norway south and east to N. Africa, Siberia, Himalayas, Israel. ksacta By streams and rivers, marshes, woods and wet fens on richer soils[17].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Branches of older wood as long as 2.5 metres can be used[1]. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta A very easily grown plant, it succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200] and dislikes poor thin soils[186]. Very tolerant of maritime exposure and atmospheric pollution[75, 186]. Trees respond well to coppicing or pollarding[186]. Best planted into its permanent position as soon as possible, trees respond badly to transplanting unless they are moved regularly. The root system is rather aggressive and can cause problems with drains[200]. A very important food plant for the caterpillars of many species of butterflies[30] and a good bee plant, providing an early source of nectar and pollen[11]. A very good wildlife habitat, more than 200 species of insects are associated with this tree[24]. There are many sub-species and cultivars in this species[182]. *S. alba caerulea* is the cricket bat willow, cultivated for its wood[11, 131]. *S. alba vitellina*. (L.)Stokes has been cultivated for its very tough stems that are used as tie rods in basket making[123, 131]. The cultivar 'Cardinal' is also grown for its use in basket making[131]. This species is used commercially in papermaking[189]. Hybridizes freely with other members of this genus, especially *S. fragilis*, to which it is closely related[11]. Trees cast a relatively light shade. Dioecious. Male and female plants must be grown if seed is required. Special Features: Attractive foliage, Not North American native, Wetlands plant, Inconspicuous flowers or blooms.

2.2.6 Populus tremula

Aspen Poplar, European aspen, Aspen
2 tlin 5 cyuc 1 kcaak 3 tcan 2 dwat 18 m hkuc
hmimhtut cyutli tucphja7n hwicli hyishrasljutnykfrip hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta Populus tremula is a deciduous Tree growing to 18 m (59ft) by 10 m (32ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 2. It is in flower from February to March, and the seeds ripen from May to June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Wind. The plant is not self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid and neutral soils. It can grow in semi-shade (light woodland) or

no shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure.

kcakta Edible Parts: Inner barkEdible Uses: Inner bark - dried, ground into a powder then added to flour and used for making bread etc[2, 115]. A famine food, it is only used when all else fails[177].

tcanta Charcoal Shelterbelt Soil conditioner WoodA very fast growing and wind resistant tree, it can be planted to provide a shelterbelt[11, 24]. Trees can also be planted to improve heavy clay soils in neglected woodlands[200]. Wood - very soft, elastic, easily split, rather woody in texture, without smell or taste, of low flammability, not durable, very resistant to abrasion. It makes a high quality paper and is also used to make a very good charcoal[11, 21, 46, 100, 115].

dwatta Anodyne Antiinflammatory Bach Diuretic Expectorant Febrifuge StimulantThe bark and the leaves are mildly diuretic, expectorant and stimulant[9, 165]. The plant is seldom used medicinally, but is sometimes included in proprietary medicines for chronic prostate and bladder disorders[9]. Although no specific mention has been seen for this species, the bark of most, if not all members of the genus contain salicin, a glycoside that probably decomposes into salicylic acid (aspirin) in the body[213, 238]. The bark is therefore anodyne, anti-inflammatory and febrifuge. It is used especially in treating rheumatism and fevers, and also to relieve the pain of menstrual cramps[238]. The plant is used in Bach flower remedies - the keywords for prescribing it are 'Vague fears of unknown origin', 'Anxiety' and 'Apprehension'[209].

nrakta Europe, including Britain, from Iceland south and east to N. Africa, temperate Asia to Japan. ksacta Found in open woodlands and scrubby heathland, usually on poorer soils where it is sometimes dominant[17].

propta Seed - must be sown as soon as it is ripe in spring[113]. Poplar seed has an extremely short period of viability and needs to be sown within a few days of ripening[200]. Surface sow or just lightly cover the seed in trays in a cold frame. Prick out the seedlings into individual pots when they are large enough to handle and grow them on in the old frame. If sufficient growth is made, it might be possible to plant them out in late summer into their permanent positions, otherwise keep them in the cold frame until the following late spring and then plant them out. Most poplar species hybridize freely with each other, so the seed may not come true unless it is collected from the wild in areas with no other poplar species growing[11]. Cuttings of mature wood in November/December in a sheltered outdoor bed. This species is rather difficult from cuttings[11, 113]. Suckers in early spring[78]. Root cuttings in the winter[200].

nwatta A very easily grown plant, it does well in a heavy cold damp soil[1], preferring a neutral to acid soil, and avoiding calcareous soils[11, 24]. The sub-species *P. tremula davidiana*. (Dode.)Schneid. is tolerant of extremely alkaline soils[200]. The species generally prefers a deep rich well-drained circumneutral soil, growing best in the south and east of Britain[11, 200]. Growth is much less on wet soils, on poor acid soils and on thin dry soils[11]. This species grows well on poor soils, probably because of its intolerance of competition[186]. Plants are very tolerant of exposure[24], doing well in cold exposed sites so long as sufficient moisture is present. Plants dislike shade[186], and are intolerant of root or branch competition[200]. A fast growing tree but it is short-lived[186]. Trees produce suckers freely and can form dense thickets[186]. The aspen is a very good wildlife plant, it has over 90 associated insect species[24] and is a food plant for the green hairstreak butterfly[186]. Poplars have very extensive and aggressive root systems that can invade and damage drainage systems. Especially when grown on clay soils, they should not be planted within 12 metres of buildings since the root system can damage the building's foundations by drying out the soil[11]. Can be planted to improve heavy clay soils in neglected woodlands[200]. Dioecious. Male and female plants must be grown if seed is required. Special Features:Not North American native, Wetlands plant, Inconspicuous flowers or blooms.

2.2.7 Salix pentandra

Bay Willow, Laurel willow

2 tlin 5 cyuc 1 kcaak 3 tcan 3 dwat 10 m hkuc
htut cyutli tucphja7n hwicli hrasljutnykfrip hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

Emetic Refrigerant Stimulant Stomachic Tonic VasodilatorThe leaves are anodyne, antiseptic, antispasmodic, astringent, carminative, cholagogue, diaphoretic, emetic, refrigerant, stimulant, stomachic, tonic and vasodilator[4, 9, 21]. A tea made from the leaves has traditionally been used in the treatment of fevers, headaches, digestive disorders and various minor ailments[222]. It is also used as a mouth-wash and a gargle for treating sore throats, ulcers, bad breath etc[9]. The leaves are harvested as the plant comes into flower and can be dried for later use[238]. The essential oil in the leaves is antiseptic, though it is toxic in large doses[222].

nrakta Europe, including Britain, south and east to N. Africa and southwest Asia Also in S. Africa, Madeira ksacta Swamps, fen, marshes, near rivers, streams and ponds, in wet woods[9].

propta Seed - sow spring in a cold frame. Germination is usually fairly quick. Prick out the seedlings into individual pots when they are large enough to handle and plant them out in the summer. *Mentha* species are very prone to hybridisation and so the seed cannot be relied on to breed true. Even without hybridisation, seedlings will not be uniform and so the content of medicinal oils etc will vary. When growing plants with a particular aroma it is best to propagate them by division[K]. Division can be easily carried out at almost any time of the year, though it is probably best done in the spring or autumn to allow the plant to establish more quickly. Virtually any part of the root is capable of growing into a new plant. Larger divisions can be planted out direct into their permanent positions. However, for maximum increase it is possible to divide the roots up into sections no more than 3cm long and pot these up in light shade in a cold frame. They will quickly become established and can be planted out in the summer.

nwatta Succeeds in most soils and situations so long as the soil is not too dry[1, 16, 200]. Grows well in heavy clay soils. A sunny position is best for production of essential oils, but it also succeeds in partial shade. Plants can grow in water up to 15cm deep[24]. Most mints have fairly aggressive spreading roots and, unless you have the space to let them roam, they need to be restrained by some means such as planting them in containers that are buried in the soil[K]. Hybridizes freely with other members of this genus. The whole plant, especially when bruised, has a pungent aroma of bergamot[245]. The flowers are especially attractive to bees and butterflies[24]. A good companion for brassicas[14]. Members of this genus are rarely if ever troubled by browsing deer[233].

5.2.16 Quercus petraea

Sessile Oak, Durmast oak

5 tlin 8 cyuc 2 kcaak 0 tcan 3 dwat 40 m hkuc
hmimhtut cyutli tucphja7n hwicli jlutnykfrip hpa7tli frusflat-prikkwis hpa7tmwatli hjonli

drutta Quercus petraea is a deciduous Tree growing to 40 m (131ft) by 25 m (82ft) at a slow rate. See above for USDA hardiness. It is hardy to UK zone 4. It is in flower from April to May, and the seeds ripen from September to October. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. It is noted for attracting wildlife. Suitable for: medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very acid soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure.

kcakta Edible Parts: SeedEdible Uses: Coffee GumSeed - cooked[2, 5, 8, 13]. It can be dried, ground into a powder and used as a thickening in stews etc or mixed with cereals for making bread. The seed contains bitter tannins, these can be leached out by thoroughly washing the seed in running water though many minerals will also be lost. Either the whole seed can be used or the seed can be dried and ground it into a powder. It can take several days or even weeks to properly leach whole seeds, one method was to wrap them in a cloth bag and place them in a stream. Leaching the powder is quicker. A simple taste test can tell when the tannin has been leached. The traditional method of preparing the seed was to bury it in boggy ground overwinter. The germinating seed was dug up in the spring when it would have lost most

of its astringency. The roasted seed is a coffee substitute[21, 61]. An edible gum is obtained from the bark[177].

tcanta Basketry Charcoal Compost Fuel Gum Ink Repellent Tannin WoodA mulch of the leaves repels slugs, grubs etc, though fresh leaves should not be used as these can inhibit plant growth[20, 201]. Oak galls are excrescences that are sometimes produced in great numbers on the tree and are caused by the activity of the larvae of different insects. The insects live inside these galls, obtaining their nutrient therein. When the insect pupates and leaves, the gall can be used as a rich source of tannin, that can also be used as a dyestuff[4]. The bark is an ingredient of 'Quick Return' herbal compost activator[32]. This is a dried and powdered mixture of several herbs that can be added to a compost heap in order to speed up bacterial activity and thus shorten the time needed to make the compost[K]. The bark is very rich in calcium[13]. An ink is made from the oak galls, mixed with salts of iron[7, 66]. The wood is a source of tar, quaiacal, acetic acid, creosote and tannin[123]. Tannin is extracted commercially from the bark and is also found in the leaves[223]. Wood - hard, tough, durable even under water. It is highly valued for furniture, construction etc[4, 13, 66]. It is also a good fuel and charcoal[6, 61]. Trees can be coppiced to provide material for basket making, fuel, construction etc[23].

dwatta Antiseptic Astringent Bach Decongestant Dysentery Haemostatic TonicThe oak tree has a long history of medicinal use. It is anti-inflammatory, antiseptic, astringent, decongestant, haemostatic and tonic[4, 7, 9, 13, 21, 165]. The bark is the part of the plant that is most commonly used[4], though other parts such as the galls, seeds and seed cups are also sometimes used[7]. A decoction of the bark is useful in the treatment of chronic diarrhoea, dysentery, intermittent fevers, haemorrhages etc[4]. Externally, it is used to bathe wounds, skin eruptions, sweaty feet, piles etc[9]. It is also used as a vaginal douche for genital inflammations and discharge, and also as a wash for throat and mouth infections[9]. The bark is harvested from branches 5 - 12 years old, and is dried for later use[9]. Any galls produced on the tree are strongly astringent and can be used in the treatment of haemorrhages, chronic diarrhoea, dysentery etc[4]. The plant is used in Bach flower remedies - the keywords for prescribing it are 'Despondency', 'Despair, but never ceasing effort'[209]. A homeopathic remedy is made from the bark. It is used in the treatment of disorders of the spleen and gall bladder[9].

nrakta Europe, including Britain, from Norway south and east to Spain, S.W. Russia and Greece. ksacta Woods, especially on acid soils and in the western part of Britain, where it is often dominant[17].

propta Seed - it quickly loses viability if it is allowed to dry out. It can be stored moist and cool overwinter but is best sown as soon as it is ripe in an outdoor seed bed, though it must be protected from mice, squirrels etc. Small quantities of seed can be sown in deep pots in a cold frame. Plants produce a deep taproot and need to be planted out into their permanent positions as soon as possible, in fact seed sown in situ will produce the best trees[11]. Trees should not be left in a nursery bed for more than 2 growing seasons without being moved or they will transplant very badly.

nwatta Landscape Uses:Specimen, Street tree. Prefers a good deep fertile loam which can be on the stiff side[11]. Dislikes heavy clay[98]. Young plants tolerate reasonable levels of shade[200]. Found mainly on acid soils in the wild. Thrives in well drained soils but is also tolerant of periodic flooding[186]. Tolerates exposure and strong winds if these are not salt-laden[186]. A very important timber tree in Britain, it is also a food plant for the caterpillars of many species of butterfly[30], there are 284 insect species associated with this tree[24]. Trees were often coppiced or pollarded in the past for their wood[186], though this is best done on a long rotation of perhaps 50 years. The tree flowers on new growth produced in spring, the seed ripening in its first year[200, 229]. Older trees have a thick corky bark and this can protect them from forest fires, young trees will often regenerate from the base if cut down or killed back by a fire[186]. Trees transplant badly unless moved regularly[11]. Hybridizes freely with other members of the genus[200]. Immune to attacks by the Tortix moth[1]. This species is notably resistant to honey fungus[88, 200]. Special Features:Not North American native, Inconspicuous flowers or blooms.

in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

5.2.14 Crataegus hupehensis

5 tlin 9 cyuc 3 kcaak 0 tcan 2 dwat 5 m hkuc hmimhtut cyuti hgattcuphja7n hwicli hrasjutnykfrp hpa7tli flatprklwsslip hpa7tmwatli xre7t hjonli

drutta Crataegus hupehensis is a deciduous Tree growing to 5 m (16ft 5in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower from May to June, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[46, 105, 177, 200]. Esteemed in China, the fruit is large but insipid[61]. The fruit is up to 2.5cm in diameter[200, 266]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many Hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta E. Asia - W. China. ksacta Thickets on slopes at elevations of 500 - 2000 metres in Henan, Hubei, Hunan, Jiangsu, Jiangxi, Shaanxi, Shanxi, Sichuan and Zhejiang Provinces[266].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. Cultivated for its edible fruit in China[46, 61, 109].

5.2.15 Mentha aquatica

Water Mint

5 tlin 9 cyuc 3 kcaak 3 tcan 3 dwat 1 m hkuc hmimhtut cyuti tcuphja7nhs6s hwicli hrasjutnykfrp hpa7tli flatprklwis hpa7tmwatli xre7t hjonli

drutta Mentha aquatica is a PERENNIAL growing to 1 m (3ft 3in) by 1 m (3ft 3in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower from July to October, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can grow in water.

pcista Although no records of toxicity have been seen for this species, large quantities of some members of this genus, especially when taken in the form of the extracted essential oil, can cause abortions so some caution is advised.

kcaakta Edible Parts: LeavesEdible Uses: Condiment TeaLeaves - raw or cooked. A strong distinctive peppermint-like fragrance[183]. Used as a flavouring in salads or cooked foods[5]. The leaves are too pungent for most people to use as a flavouring[244]. A herb tea is made from the leaves[21, 183].

tcanta Repellent StrewingThe plant repels flies, mice and rats[14, 20, 244]. It has a pleasant, fresh scent and was formerly used as a strewing herb and has been strewn in granaries to keep mice and rats off the grain[14, 244]. The plant, harvested before flowering, yields about 0.8% essential oil[240]. The fresh or dried plant is very good when used in herbal baths and can also be used in herb pillows[244]. Water garden.

dwatta Anodyne Antidiarrhoeal Antiseptic Antispasmodic Astringent Carminative Cholagogue Diaphoretic

drutta Salix pentandra is a deciduous Tree growing to 10 m (32ft) by 10 m (32ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen in June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required). and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil.

pcista Gastrointestinal bleeding and kidney damage possible. Avoid concurrent administration with other aspirin-like drugs. Avoid during pregnancy. Drug interactions associated with salicylates applicable [301].

kcaakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - cooked. Not very palatable[172].

tcanta Basketry Pot-pourriLandscape Uses: Specimen. Special Features: Not North American native, Inconspicuous flowers or blooms. The stems are very flexible and are used in basket making[131]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. The dried leaves have a pleasant aromatic aroma and can be used in pot-pourri[245]. Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11].

dwatta Anodyne Dysentery FebrifugeThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226]. The bark of this species is used interchangeably with *S. alba*. It is taken internally in the treatment of rheumatism, arthritis, gout, inflammatory stages of autoimmune diseases, diarrhoea, dysentery, feverish illnesses, neuralgia and headache[238]. The bark is removed during the summer and dried for later use[238]. The leaves are used internally in the treatment of minor feverish illnesses and colic[238]. The leaves can be harvested throughout the growing season and are used fresh or dried[238]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve *Salix* / Willow for diseases accompanied by fever, rheumatic ailments, headaches (see [302] for critics of commission E).

nrakta Europe, including Britain, from Norway south and east to the Pyrenees, Siberia, Caucasus, W. Asia. ksacta Stream-sides, marshes, fens and wet woods, ascending to 450 metres[17]. Native in N. Britain, planted elsewhere[17].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. A very ornamental plant[1]. Best planted into its permanent position when young[11]. The root system is rather aggressive and can cause problems with drains[200]. The dried or bruised leaves have a sweet aromatic fragrance[245]. The scent resembles oil of wintergreen and is due to the presence of salicyl aldehyde[245]. Cultivated for its use in basket making, there are several named varieties[131]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

3 3 tlin

3.0.1 Viburnum opulus

Guelder Rose, Cramp Bark, European cranberrybush, American cranberrybush, Crampbark, European Highh 3 tlin 8 cyuc 3 kcaak 3 tcan 3 dwat 5 m hkuc hmimhtut cyuti tcuphja7n hwicli hrasjutnykfrp hpa7tli flatprklwsslip hpa7tmwatli fyifxre7t hjonli

drutta Viburnum opulus is a deciduous Shrub growing to 5 m (16ft) by 5 m (16ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 3 and is not frost tender. It is in flower from June to July, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Insects. The plant is self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista Large quantities of the fruit can cause vomiting and diarrhoea[10, 65]. The fruit is of very low or zero toxicity, it only causes mild upsets when eaten unripe or in large quantities[65, 76].

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[2, 3, 5, 46]. The fruit is up to 8.5mm in diameter but with a large seed[200]. A sour taste, it is best cooked. The crushed fruit has an unpleasant smell[4]. Used as a cranberry substitute in making, jellies, preserves etc[183]. Some caution is advised, see notes on toxicity at top of the page.

tcanta Dye Hedge Hedge Ink WoodA red dye is obtained from the fruit[13]. An ink can be made from the dried berries[4]. Plants can be grown as a tall hedge[29], they are rather bare in winter though[K]. The wood can be used to make skewers[4].

dwatta Antispasmodic Astringent Birthing aid Homeopathy SedativeGuelder rose is a powerful antispasmodic and is much used in the treatment of asthma, cramps and other conditions such as colic or painful menstruation[254]. It is also used as a sedative remedy for nervous conditions[254]. The bark is antispasmodic, astringent and sedative[4, 9, 46, 165, 213]. The bark contains 'scoletin', a coumarin that has a sedative effect on the uterus[238]. A tea is used internally to relieve all types of spasms, including menstrual cramps, spasms after childbirth and threatened miscarriage[9, 222, 238]. It is also used in the treatment of nervous complaints and debility[4, 46, 165, 213]. The bark is harvested in the autumn before the leaves change colour, or in the spring before the leaf buds open. It is dried for later use[238]. The leaves and fruits are antiscorbatic, emetic and laxative[4, 222]. A homeopathic remedy is made from the fresh bark[9]. It is used in the treatment of menstrual pain and spasms after childbirth[9].

nrakta Europe, including Britain, from Scandinavia south and east to Spain, north and west Asia. ksacta Hedges, scrub and woodland, usually on damp soils[3, 13, 17].

propta Seed - best sown in a cold frame as soon as it is ripe. Germination can be slow, sometimes taking more than 18 months. If the seed is harvested 'green' (when it has fully developed but before it has fully ripened) and sown immediately in a cold frame, it should germinate in the spring[80]. Stored seed will require 2 months warm then 3 months cold stratification and can still take 18 months to germinate[113]. Prick out the seedlings into individual pots when they are large enough to handle and grow them on in a cold frame or greenhouse. Plant out into their permanent positions in late spring or early summer of the following year. Cuttings of soft-wood, early summer in a frame[200]. Pot up into individual pots once they start to root and plant them out in late spring or early summer of the following year. Cuttings of half-ripe wood, 5 - 8 cm long with a heel if possible, July/August in a frame[78, 113]. Plant them into individual pots as soon as they start to root. These cuttings can be difficult to overwinter, it is best to keep them in a greenhouse or cold frame until the following spring before planting them out[113]. Cuttings of mature wood, winter in a frame. They should root in early spring - pot them up when large enough to handle and plant them out in the summer if sufficient new growth is made, otherwise keep them in a cold frame for the next winter and then plant them out in the

spring. Layering of current seasons growth in July/August. Takes 15 months[78].

nwatta Landscape Uses:Border, Massing, Screen, Specimen. An easily grown plant, it succeeds in most soils but is ill-adapted for poor soils and for dry situations[1]. It prefers a deep rich moist loamy soil in a sunny position[11]. Succeeds in semi-shade but does not grow or fruit so well in such a position[186]. Grows well in heavy clay soils and on chalk[184]. Does not do well on very acid soils. Best if given shade from the early morning sun in spring[200]. A very ornamental plant[1], it is hardy to about -30°c[184] and is often grown in the flower garden. There are many named varieties[184]. Guelder rose regenerates quickly if it is cut to the ground, it can also produce suckers and will often form thickets[186]. The plant is an alternative host for the broad bean aphid[11]. Special Features: Not North American native, Naturalizing, Attractive flowers or blooms. In garden design, as well as the above-ground architecture of a plant, root structure considerations help in choosing plants that work together for their optimal soil requirements including nutrients and water. The root pattern is fleshy. Thick or swollen - fibrous or tap root [2-1].

3.0.2 Rosa canina

Dog Rose

3 tlin 7 cyuc 3 kcaak 4 tcan 3 dwat 3 m hkuc hmimhtut cyutli tucphja7n hwicli hrasjutnykfrrip hpa7tli flatprikwis hpa7tmwatli fyifxre7t hjonli

drutta Rosa canina is a deciduous Shrub growing to 3 m (9ft) by 3 m (9ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 3. It is in flower from June to July, and the seeds ripen from October to December. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies, beetles, Lepidoptera (Moths and Butterflies), Apomictic (reproduce by seeds formed without sexual fusion). The plant is self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils, prefers well-drained soil and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure.

pista There is a layer of hairs around the seeds just beneath the flesh of the fruit. These hairs can cause irritation to the mouth and digestive tract if ingested.

kcaakta Edible Parts: Flowers Fruit SeedEdible Uses: Coffee TeaFruit - raw or cooked[2, 4, 13]. It can be used in making delicious jams, syrups etc[7]. The syrup is used as a nutritional supplement, especially for babies[238]. The fruit can also be dried and used as a tea[9, 12, 183]. Frost softens and sweetens the flesh[12]. The fruit is up to 30mm in diameter[200], but there is only a thin layer of flesh surrounding the many seeds[K]. Some care has to be taken when eating this fruit, see the notes above on known hazards. The seed is a good source of vitamin E[102], it can be ground and mixed with flour or added to other foods as a supplement. Be sure to remove the seed hairs[102]. The dried leaves are used as a tea substitute[46, 61, 183]. A coffee substitute according to another report[7]. Petals - raw or cooked[177]. The base of the petal may be bitter so is best removed[183]. Eaten as a vegetable in China[66]. The petals are also used to make an unusual scented jam[7].

tcanta Hedge HedgePlants make a dense and stock-proof hedge, especially when trimmed[186].

dwatta Antirheumatic Astringent Bach Cancer Carmine Diuretic Laxative Ophthalmic Tonic Vermifuge Vitamin CThe petals, hips and galls are astringent, carminative, diuretic, laxative, ophthalmic and tonic[7, 9, 14, 21, 165]. The hips are taken internally in the treatment of colds, influenza, minor infectious diseases, scurvy, diarrhoea and gastritis[238]. A syrup made from the hips is used as a pleasant flavouring in medicines and is added to cough mixtures[238]. A distilled water made from the plant is slightly astringent and is used as a lotion for delicate skins[7]. The seeds have been used as a vermifuge[7]. The plant is used in Bach flower remedies - the keywords for prescribing it are 'Resignation' and 'Apathy'[209]. The fruit of many members of this genus is a very rich source of vitamins and minerals, especially in vitamins A, C and E, flavanoids and other bioactive compounds. It is also a fairly good source of essential

fatty acids, which is fairly unusual for a fruit. It is being investigated as a food that is capable of reducing the incidence of cancer and also as a means of halting or reversing the growth of cancers[214]. Ascorbic acid in Dog Rose shells (vitamin C, 0.2 to 2.4%).

nrakta Europe, including Britain, from Norway south and east to N. Africa and southwest Asia. ksacta Hedges, scrub, woods, roadsides, banks etc[9].

propta Seed. Rose seed often takes two years to germinate. This is because it may need a warm spell of weather after a cold spell in order to mature the embryo and reduce the seedcoat[80]. One possible way to reduce this time is to scarify the seed and then place it for 2 - 3 weeks in damp peat at a temperature of 27 - 32°c (by which time the seed should have imbibed). It is then kept at 3°c for the next 4 months by which time it should be starting to germinate[80]. Alternatively, it is possible that seed harvested 'green' (when it is fully developed but before it has dried on the plant) and sown immediately will germinate in the late winter. This method has not as yet(1988) been fully tested[80]. Seed sown as soon as it is ripe in a cold frame sometimes germinates in spring though it may take 18 months. Stored seed can be sown as early in the year as possible and stratified for 6 weeks at 5°c[200]. It may take 2 years to germinate[200]. Prick out the seedlings into individual pots when they are large enough to handle. Plant out in the summer if the plants are more than 25cm tall, otherwise grow on in a cold frame for the winter and plant out in late spring. Cuttings of half-ripe wood with a heel, July in a shaded frame. Overwinter the plants in the frame and plant out in late spring[78]. High percentage[78]. Cuttings of mature wood of the current seasons growth. Select pencil thick shoots in early autumn that are about 20 - 25cm long and plant them in a sheltered position outdoors or in a cold frame[78, 200]. The cuttings can take 12 months to establish but a high percentage of them normally succeed[78]. Division of suckers in the dormant season. Plant them out direct into their permanent positions. Layering. Takes 12 months[11].

nwatta Succeeds in most soils[11, 200]. Grows well in heavy clay soils. Prefers a circumneutral soil and a sunny position with its roots in the shade[11, 200]. When grown in deep shade it usually fails to flower and fruit[186]. Succeeds in wet soils but dislikes water-logged soils or very dry sites[186]. Tolerates maritime exposure[186]. The fruit attracts many species of birds, several gull wasps and other insects use the plant as a host[186] A very polymorphic species, it is divided into a great number of closely related species by some botanists[1, 17]. The leaves, when bruised, have a delicious fragrance[245]. The flowers are also fragrant. Grows well with alliums, parsley, mugwort and lupins[18, 20]. Garlic planted nearby can help protect the plant from disease and insect predation[18, 20]. Grows badly with boxwood[18]. Hybridizes freely with other members of this genus[80]. Plants in this genus are notably susceptible to honey fungus[200].

3.0.3 Asclepias incarnata

Swamp Milkweed, Swamp Butterfly Weed, Marsh Milkweed 3 tlin 8 cyuc 3 kcaak 3 tcan 2 dwat 1.2 m hkuc hmimhtut cyutli kyuctuphja7n hwicli hrasjutnykfrrip hpa7tli flatprikwis hpa7tmwatli fyifxre7t hjonli

drutta Asclepias incarnata is a PERENNIAL growing to 1.2 m (4ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 3. It is in flower from July to August, and the seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, insects, Lepidoptera (Moths and Butterflies). The plant is self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy) and medium (loamy) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil.

pista Although no specific reports have been seen for this species, many, if not all, members of this genus contain toxic resinoids, alkaloids and cardiac glycosides[274]. They are usually avoided by grazing animals[274]. The leaves and the stems might be poisonous[20].

kcaakta Edible Parts: Flowers Leaves Oil SeedpodEdible Uses: Oil SweetenerUnopened flower buds - cooked[46, 61, 161]. Tasting somewhat like peas[85]. They can also be

pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta Industrial Crop: Medicinal Management: Standard Minor Global CropA very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Most trees have very few thorns, some forms are thorn-free[202]. A very ornamental plant[1, 11]. The sub-species C. pinnatifida major has larger fruit than the type species, it is cultivated for this fruit in China[46, 178, 200] Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. For polyculture design as well as the above-ground architecture (form - tree, shrub etc. and size shown above) information on the habit and root pattern is also useful and given here if available. The plant growth habit is a standard with a non-suckering single trunk [1-2]. The root pattern is a tap root similar to a carrot going directly down [1-2].

5.2.12 Crataegus pentagyna

5 tlin 9 cyuc 3 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjutnykfrrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus pentagyna is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower in June. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[2, 46, 105]. It can also be made into preserves[61] and can be dried for later use. The fruit is about 1cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta S.E. Europe to the Caucasus. ksacta Woodland margins[45, 50].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold

stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Cultivated in China for its edible fruit, it is sold in the local markets[46, 61]. This report might be a mistake, probably applying instead to C. pinnatifida[K] Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

5.2.13 Crataegus aprica

5 tlin 9 cyuc 3 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgatkyuctuphja7n hwicli hrasjutnykfrrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus aprica is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower in May, and the seeds ripen in November. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. A good dessert fruit, it is sweet and rather juicy[82]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is about 12mm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta South-eastern N. America - Virginia to Tennessee and Georgia. ksacta Dry woods in the foothills of the Appalachians, 450 - 1000 metres[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate

transplanted[11].

5.2.10 Crataegus flava

Summer Haw, Yellowleaf hawthorn
5 tin 9 cyuc 3 kcaak 3 tcan 2 dwat 8 m hkuc
hmimhtut cyutli hgatkycutcuphja7n hwicli hrasjlutnyikfri
p hpa7tli flatprikliwsslip hpa7tmwatli xre7t hjonli

drutta Crataegus flava is a deciduous Shrub growing to 8 m (26ft) by 10 m (32ft). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

ckaakta Edible Parts: Fruit(Edible Uses: Fruit - raw or cooked[46, 61, 95, 161]. well-flavoured[2]. Dry and mealy[82], it is highly esteemed in N. America as a wild fruit gathered for making jellies[183]. The yellow fruit is pear-shaped[245], it is up to 16mm in diameter and is borne in small clusters[229]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta South-eastern N. America - Virginia to Florida, west to Mississippi. ksacta Open woods and low hills in dry sandy or gravelly soils[43, 82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open,

the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. The fruit is sparsely produced in Britain[11]. A tree at Kew in September 1993 was fruiting heavily[K].

5.2.11 Crataegus pinnatifida

Chinese Haw

5 tin 9 cyuc 3 kcaak 2 tcan 3 dwat 7 m hkuc
hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnyikfri
p hpa7tli flatprikliwsslip hpa7tmwatli xre7t hjonli

drutta Crataegus pinnatifida is a deciduous Tree growing to 7 m (23ft) by 7 m (23ft). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower in May, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

ckaakta Edible Parts: Fruit(Edible Uses: Fruit - raw or cooked[105, 177]. A pleasant flavour[178]. The orange fruit has a mealy texture with an acid taste and a slight bitterness but is fairly nice raw[K]. It ripens fairly late, specimens seen at the end of October 1998 were not quite fully ripe[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is about 15mm in diameter[200], though in some forms it can be 25mm in diameter[266]. A nutritional analysis is available[218]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Alterative Antibacterial Anticholesterolemic Antidote Antiscorbutic Cardiotonic Deobstruent Hypotensive Laxative Stimulant Stomachic VasodilatorThe fruit is antibacterial, anticholesterolemic, cardiotonic, hypotensive, stomachic, vasodilator. It contains several medically active constituents including flavonoids and organic acids[279]. It has been shown to lower blood cholesterol levels and to improve blood circulation[176, 238, 279]. It is used in Korea in the treatment of abdominal distension, pain, diarrhoea and to induce menstruation[279]. The dried fruit is alterative, antiscorbutic, deobstruent, laxative, stimulant and stomachic[218]. The fruits and flowers of hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222]. The seed is recommended in the treatment of difficult labour, hernia and swollen genitals[218]. The leaves and twigs are used as an antidote to varnish poisoning (from certain Rhus spp)[218]. The root is used in the treatment of nausea and vomiting[218].

nrakta E. Asia - N. China, Korea. ksacta Riverbanks, on sandy soils or stony slopes[74]. Among shrubs, on slopes at elevations of 100 - 2000 metres[266].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to

dried and stored for later use[183]. Young shoots - cooked. An asparagus substitute[85, 106]. Tips of older shoots are cooked like spinach[85]. Young seed pods, harvested when 3 - 4 cm long - cooked. A pea-like flavour, they are very appetizing[85]. The flower clusters can be boiled down to make a sugary syrup[85].

tcanta Fibre Latex Oil Pollution Stuffing WaxA good quality fibre is obtained from the bark[46, 57, 61, 95, 112, 169]. It is used in twine, cloth etc[112]. It is easily harvested in late autumn, after the plants have died down, by simply pulling it off the dead stems[112]. The seed floss is used to stuff pillows etc or is mixed with other fibres to make cloth[57, 171]. It is a Kapok substitute, it is used in Life Jackets or as a stuffing material[169, 171]. It is very water repellent. The floss has also been used to mop up oil spills at sea. Rubber can be made from latex contained in the leaves and stems[57]. Pods contain an oil and a wax which are of potential importance[171]. The dogbane-milkweed family Asclepias, Apocynum, Calotropis, and Trachomitum spp) has been used for fiber industrial crops for millennia with a number in cultivation as regional crops. All of these crops are dual-purpose fibres, offering bast fibres from the stem and seed finer or 'floss' in the fruit pods. Many have also been identified as potential hydrocarbon crops due to high latex content. Could be integrated into various agroforestry systems rather than as monocultures [1-1].

dwatta Anthelmintic Carminative Diuretic Emetic Laxative Stomachic A tea made from the roots is anthelmintic, carminative, diuretic, emetic, strongly laxative and stomachic[4, 61, 222, 257]. The tea is said to remove tapeworms from the body in one hour[257]. It has also been used in the treatment of asthma, rheumatism, syphilis, worms and as a heart tonic[4, 207, 222]. An infusion of the roots is used as a strengthening bath for children and adults[257].

nrakta N. America - Quebec to Manitoba and Wyoming, south to Texas and New Mexico. ksacta Swamps, wet thickets and shores[43].

propta Seed - best sown in a greenhouse as soon as it is ripe in the autumn or in late winter[134, 169]. We have also had good results from sowing the seed in the greenhouse in early spring[K], though stored seed might need 2 - 3 weeks cold stratification[134]. Germination usually takes place in 1 - 3 months at 18°C[134]. As soon as the seedlings are large enough to handle, prick them out into individual pots and grow them on in the greenhouse for their first winter. Plant out when they are in active growth in late spring or early summer and give them some protection from slugs until they are growing away strongly. Division in spring. With great care since the plant resents root disturbance. Pot the divisions up and place them in a lightly shaded position in the greenhouse until they are growing away strongly, then plant them out in the summer, giving them some protection from slugs until they are established.. Basal cuttings in late spring. Use shoots about 10cm long with as much of their white underground stem as possible. Pot them up individually and place them in a lightly shaded position in a greenhouse until they are rooting and growing actively. If the plants grow sufficiently, they can be put into their permanent positions in the summer, otherwise keep them in the greenhouse until the following spring and when they are in active growth plant them out into their permanent positions. Give them some protection from slugs until they are established.

nwatta Experimental Crop Industrial Crop: Hydrocarbon Management: HayLandscape Uses:Border, Foundation, Massing. Prefers a well-drained light rich or peaty soil[1, 200]. Requires a moist soil and a sunny position, doing well by water[111, 134]. Succeeds on dry soils and on all soil types[112]. Plants are hardy to at least -25°C[187]. A very ornamental plant[1], the flowers are very attractive to butterflies[169]. The flower of many members of this genus can trap insects between its anther cells, the struggles of the insect in escaping ensure the pollination of the plant[207]. Many members of this genus seem to be particularly prone to damage by slugs. The young growth in spring is especially vulnerable, but older growth is also attacked and even well-established plants have been destroyed in wet years[K]. Plants resent root disturbance and are best planted into their final positions whilst small[134]. Special Features:Attractive foliage, North American native, Naturalizing, Wetlands plant, Attracts butterflies. Suitable for cut flowers, Fragrant flowers. The plant is heat tolerant in zones 9 through 2. (Plant Hardiness Zones show how well plants

withstand cold winter temperatures. Plant Heat Zones show when plants would start suffering from the heat. The Plant Heat Zone map is based on the number of "heat days" experienced in a given area where the temperature climbs to over 86 degrees F (30°C). At this temperature, many plants begin to suffer physiological damage. Heat Zones range from 1 (no heat days) to 12 (210 or more heat days). For example Heat Zone. 11-1 indicates that the plant is heat tolerant in zones 11 through 1.) For polyculture design as well as the above-ground architecture (form - tree, shrub etc. and size shown above) information on the habit and root pattern is also useful and given here if available. The plant growth habit is a clumper with limited spread [1-2]. The root pattern is rhizomatous with underground stems sending roots and shoots along their length [1-2].

3.0.4 Filipendula ulmaria

Meadowsweet, Queen of the meadow, Double Lady of the Meadow, European Meadowsweet
3 tin 9 cyuc 3 kcaak 3 tcan 3 dwat 1.2 m hkuc
hmimhtut cyutli tcuphja7n hwicli hrasjlutnyikfri
p hpa7tli prikliw hpa7tmwatli fyfxre7t hjonli

drutta Filipendula ulmaria is a PERENNIAL growing to 1.2 m (4ft) by 0.4 m (1ft 4in) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 2 and is not frost tender. It is in flower from June to August, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies, beetles. The plant is self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista Avoid for asthmatics. Use by children for diarrhoea not recommended or for children under 12 due to salicylate content (risk of Reye's syndrome) [301].

ckaakta Edible Parts: Flowers Leaves Root(Edible Uses: Condiment TeaAll parts of the plant are edible. Root - cooked[2, 141]. Young leaves - cooked as a flavouring in soups[177]. Young leaves, flowers and roots are brewed into a tea[183]. The dried leaves are used as a flavouring[12, 100], especially as a sweetener in herb teas[13, 183]. The flowers are used as a flavouring in various alcoholic beverages and in stewed fruits[183]. Adding them to wine or beer is said to make a very healthy brew[244]. They are also made into a syrup which can be used in cooling drinks and fruit salads[183].

tcanta Dye Essential Pot-pourri StrewingA black dye is obtained from the roots. It is brown[141]. A yellow dye is obtained from the plant tops[106]. An essential oil obtained from the flower buds is used in perfumery[46, 61]. The whole plant, but especially the leaves[245], was formerly used as a strewing herb, imparting an almond-like fragrance[4, 66]. Strongly aromatic, its delightful perfume would completely fill the room[244]. Both flowers and leaves have been used in pot-pourri, retaining their scent for several months. The scent of the dried flowers becoming more and more pleasant with age[245].

dwatta Alterative Antiinflammatory Antirheumatic Antiseptic Aromatic Astringent Diaphoretic Diuretic Dysentery Homeopathy Stomachic TonicMeadowsweet has a very long history of herbal use, it was one of the three most sacred herbs of the Druids[238]. The leaves and flowering stems are alterative, anti-inflammatory, antiseptic, aromatic, astringent, diaphoretic, diuretic, stomachic and tonic[4, 9, 21, 165]. The plant is harvested in July when it is in flower and can be dried for later use[4]. The flower head contains salicylic acid, from which the drug aspirin can be synthesised[200, 238]. Unlike the extracted aspirin, which can cause gastric ulceration at high doses, the combination of constituents in meadowsweet act to protect the inner lining of the stomach and intestines whilst still providing the anti-inflammatory benefits of aspirin[254]. The herb is a valuable medicine in the treatment of diarrhoea, indeed it is considered almost specific in the treatment of children's diarrhoea[4]. It is also considered to be a useful stomachic, being used to treat hyperacidity, heartburn, gastritis and peptic ulcers, for which it is one of the most effective plant remedies[4, 238]. It is also frequently used in the treatment

of afflictions of the blood[4]. Meadowsweet is also effective against the organisms causing diphtheria, dysentery and pneumonia[238]. This remedy should not be given to people who are hypersensitive to aspirin[238]. A strong decoction of the boiled root is said to be effective, when used externally, in the treatment of sores and ulcers[244]. A homeopathic remedy is made from the fresh root[9]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve *Filipendula ulmaria* Meadowsweet for cough, bronchitis, fever and cold (see [302] for critics of commission E).

nrakta Europe, including Britain, from Iceland south and east to Spain, temperate Asia and Mongolia. ksacta Wet ground in swamps, marshes, fens, wet woods and meadows, wet rock ledges and by rivers, but not on acid peats[17].

propta Seed - best sown in the autumn in a cold frame[11]. The seed can also be sown in a cold frame in spring, germinating best at a temperature of 10 - 13°C[200]. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer if they have grown enough. If not, keep them in a cold frame for the winter and plant them out in late spring. Division in autumn or winter[200]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring.

nwatta Landscape Uses:Border, Massing, Rock garden, Seashore, Specimen, Woodland garden. Requires a humus-rich moist soil in semi-shade[200]. Succeeds in full sun only if the soil is reliably moist throughout the growing season[200]. Dislikes dry or acid soils[1, 17]. Does well in marshy soils[24, 200]. Grows well in heavy clay soils. The flowers have a strong sweet smell[4], which for many people is sickly[245]. The leaves are also aromatic[187], though the scent is very different from the flowers[4]. The leaves are pleasantly aromatic[245]. A good bee plant[24, 30]. Members of this genus are rarely if ever troubled by browsing deer[233]. Special Features:Attractive foliage, Edible, Fragrant foliage, Not North American native, Naturalizing, Suitable for cut flowers, Suitable for dried flowers, Fragrant flowers.

3.0.5 Thuja occidentalis

American Arbor-Vitae, Arborvitae, Eastern Arborvitae, Siberian Arborvitae, Northern White Cedar, Wn 3 lin 7 cyuc 2 kcaak 0 tcan 3 dwat 15 m hkuc hmimhtut cyullit cuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprklriwswilip hpa7tmwathi hjonli

drutta Thuja occidentalis is an evergreen Tree growing to 15 m (49ft) by 5 m (16ft) at a slow rate. See above for USDA hardiness. It is hardy to UK zone 2. It is in leaf all year, in flower from April to May, and the seeds ripen from September to October. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista An essential oil from the leaves is poisonous if taken in large doses[7, 65]. Nausea, vomiting, flatulence, indigestion reported. Possible asthma and seizures[301]. This plant should not be used by pregnant women[165]. Use oil internally in small quantities and under medical supervision. Death and convulsions reported with improper internal use [301].

kcakta Edible Parts: StemEdible Uses: TeaPith of young shoots - cooked[105]. It can be added to soups[177]. Pleasantly sweet, the pith was used as the basis of the soup according to one report[183]. Inner bark - cooked. It is only used in times of emergency or scarcity[213]. The inner bark can be dried and ground into a powder, then used with wheat or other cereals in making bread, biscuits etc. The leafy branchlets are used as a tea substitute[159, 177, 257] but are probably best avoided by pregnant women[165]. An aromatic flavour[183]. Another report says that the foliage and bark are used, the resulting tea is a good source of vitamin C[226].

tcanta Broom Essential Fibre Incense Repellent Tannin WoodTolerant of regular trimming, though not into the old wood, it can be grown as a hedge[226]. The fresh branches

are used as besoms[4]. Their aromatic smell serves to deodorize the house whilst sweeping[226]. The leaves have been kept in the clothes cupboard as a perfume, incense and insect repellent[257]. The leaves and stems have been used as an incense[257]. An essential oil is obtained from the leaves and branches, it is used in perfumery and in medicines[46, 57, 61, 226]. It is poisonous if taken internally[65]. This essential oil also has insect repellent properties[106]. The tough and stringy bark has been used to weave fibre bags[257]. The bark is a source of tannin[257]. Wood - light, soft, not strong, brittle, coarse grained, very durable, easily worked, does not warp[61, 82, 171, 226, 235]. It weighs 20lb per cubic foot[235]. Used especially where contact with water cannot be avoided, for canoes, garden buildings, shingles, posts etc[11, 61, 82, 171, 226].

dwatta Alterative Anthelmintic Antiinflammatory Antirheumatic Antiseptic Antiviral Aromatic Astringent Diaphoretic Diuretic Emmenagogue Homeopathy Rubefacient Vitamin C American arbor-vitae was much used by many native North American Indian tribes as a medicine to treat fevers, coughs, headaches, swollen hands and rheumatic problems[254, 257]. The plant has an established antiviral activity and is most commonly used in modern herbalism to treat warts and polyps, being prescribed both internally and externally for these conditions[254]. The plant can be used to induce menstruation and so should not be prescribed for pregnant women[238]. The recently dried leafy young twigs are alterative, anthelmintic, anti-inflammatory, antiseptic, aromatic, astringent, diaphoretic, diuretic and emmenagogue[4, 7, 21, 165]. The plant is being used internally in the treatment of cancer[238], especially cancer of the uterus[254]. A tea made from the leaves is used in the treatment for bronchitis and other respiratory problems, colds, headaches and as a cough syrup[222, 254]. The plants diuretic properties make it useful in treating acute cystitis and bed-wetting in children[254]. The leaves are used in steam baths in the treatment of rheumatism, arthritis, colds etc[222]. Externally, the leaves are used as a wash for swollen feet and burns[222]. Extracts of the leaves can be painted on painful joints or muscles as a counterirritant, improving local blood supply and thus facilitating the removal of toxins, easing pain and stiffness[254]. A tincture of the leaves has been used in the treatment of warts, piles, bed sores and fungal infections[222]. The leaves and young twigs can be harvested as required and used fresh or dried[238]. 'Oil of white cedar', obtained from the leaves, is an essential oil that is antiseptic, expectorant and rubefacient[213, 222]. It is used internally to promote menstruation and relieve rheumatism[213]. This volatile oil is toxic and poisoning from overdoses has occurred[213], it should only be used under the supervision of a qualified practitioner and should not be prescribed for pregnant women[238]. The oil also stimulates the heart and causes convulsions in high doses[213]. A tea of the inner bark is used to promote menstruation[213] and in the treatment of consumption and coughs[222]. A homeopathic remedy is made from the leaves and twigs, gathered when the tree is flowering[232]. It is used in the household as a treatment against warts, but also has a range of other applications that should only be prescribed by a competent homeopath[7, 232].

nrakta Eastern N. America - Nova Scotia to Georgia, west to Illinois and Minnesota. ksacta Swamps and cool rocky banks of streams[43], forming dense almost impenetrable forests[82, 165]. Also found on rocky soils[226]. Plants are usually found on neutral to alkaline limestone soils[226].

propta Seed - best sown when ripe in the autumn in a cold frame[113]. Stored seed germinates best if given a short cold stratification[113]. It can be sown in a cold frame in late winter. When they are large enough to handle, prick the seedlings out into individual pots and grow them on in the greenhouse for their first winter. Plant them out into their permanent positions in late spring or early summer, after the last expected frosts. If growing large quantities of plants, the seed can be sown in an outdoor seed bed in mid spring[78]. Grow the plants on for two years and then plant them out into their permanent positions in late autumn or early spring. Cuttings of half-ripe wood, 5 - 8cm with a heel, July/August in a shaded frame. Forms roots by the end of September but it should be overwintered in a frame[78]. Cuttings of almost ripe wood, 5 - 10cm with a heel, September in a cold frame. Forms roots in the following summer. Plant out in autumn or

in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[2, 105]. A pleasant acid taste[74]. The fruit is about 1.5cm in diameter[200]. A nice taste with a very good flesh to seed ratio, it can be eaten in quantity for dessert[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta WoodWood - heavy, close-grained, hard and tough. Used for tool handles etc[82, 74].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta E. Asia. Occasionally self-sows in Britain[17]. ksacta Mountain thickets and rocky slopes[50].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. A very hardy species, tolerating temperatures down to at least -18°C[202]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

5.2.9 Crataegus cuneata

Sanzashi, Chinese hawthorn 5 lin 9 cyuc 3 kcaak 3 tcan 3 dwat 15 m hkuc hmimhtut cyullit hgattcuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprklriwswilip hpa7tmwathi xre7t hjonli

drutta Crataegus cuneata is a deciduous Shrub growing to 15 m (49ft 3in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower from May

to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked and used in pies, preserves etc[147, 183]. It can also be dried for later use[177, 183]. A pleasant flavour, it is sold in local markets in China and Japan[178]. The fruit contains about 0.44% protein, 1% fat, 22.1% carbohydrate, 0.8% ash, it is rich in vitamin C, fruit acids and pectin[179]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Anodyne Anticholesterolemic Astringent Blood tonic Cardiotonic Dysentery Haemostatic Hypotensive Stomachic Vitamin CThe fruits and flowers of hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222]. The fruit is anodyne, anticholesterolemic, antiarrhetic, antidiabetic, astringent, blood tonic, cardiotonic, haemostatic and stomachic[61, 147, 176, 218]. It is used in the treatment of dyspepsia, stagnation of fatty food, abdominal fullness, retention of lochia, amenorrhoea, postpartum abdominal pain, hypertension and coronary heart disease[176].

nrakta E. Asia - China, Japan. ksacta Sunny places in upland wilds[147]. Valleys, thickets and hills at elevations of 200 - 2000 metres[266].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Although perfectly cold-hardy in most of Britain when dormant, the young growth of this species might be susceptible to spring frosts[11]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Anticholesterolemic Cardiotonic Hypotensive Stomachic VasodilatorThe fruits and flowers of hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222]. The fruits are anticholesterolemic, cardiotonic, stomachic and vasodilator. Their use improves the blood circulation and lowers blood pressure[176]. The fruits are used in the treatment of dyspepsia, stagnation of fatty foods, abdominal fullness, retention of lochia, amenorrhoea, postpartum abdominal pain, hypertension and coronary heart disease[176].

nrakta E. Asia - China. ksacta Riverbanks, on sandy soils or stony slopes[74].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A very ornamental plant, this sub-species is widely cultivated for its edible fruit in China[266], it is larger than the type species[46, 178, 200].

5.2.7 Crataegus champlainensis

Quebec hawthorn

5 tlin 9 cyuc 4 kcaak 0 tcan 2 dwat 6 m hkucc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikklwis hpa7tmwatli xre7t hjonli

drutta Crataegus champlainensis is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought.

The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. A nice sweet flavour; the fruit is about 1.5cm in diameter[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Northern N. America - Quebec to New York and Ontario. ksacta Thickets, streambanks and hillsides[204]. Limestone ridges[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. Closely related to *C. submollis*[200], and included in that species by some botanists[82]. This species is possibly no more than a part of *C. rotundifolia*[204].

5.2.8 Crataegus laciniata

5 tlin 9 cyuc 4 kcaak 0 tcan 2 dwat 6 m hkucc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikklwis hpa7tmwatli xre7t hjonli

drutta Crataegus laciniata is a deciduous Tree growing to 6 m (19ft) by 6 m (19ft). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower in June, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow

spring[78].

nwatta Landscape Uses:Firewood, Hedge, Screen, Superior hedge, Specimen. Prefers a permanently moist soil[1], it is intolerant of dry soils[11]. A useful plant for very poorly-drained soils[185, 200]. The best stands in America are on well-drained soils[11]. Grows well in heavy clay soils. Succeeds in sandy soils[188] and on chalk[200]. A very hardy tree, tolerating very cold winters with temperatures down to -46°c[200, 238]. In cold weather the leaves turn brown, becoming green again in the spring[226]. It is usually slow growing and short-lived in cultivation in Britain and rarely looks thrifty[81, 185]. However, there are some good specimens in western Britain[4, 14]. Some cultivars are more healthy, 'Lutea' is growing very well in several places and 'Spiralis' is also growing well[185]. Trees live 200 - 300 years in the wild[226]. Sometimes planted as a timber tree in C. Europe[50]. Plants cannot regenerate from old wood. Pruning is not normally necessary for this species, any pruning that is carried out should be done with care[238]. The wood and the foliage are strongly aromatic[226]. The crushed leaves have a scent of apples[185]. Special Features: Attracts birds, North American native, Wetlands plant, Inconspicuous flowers or blooms.

3.0.6 Filipendula kamschatica

3 tlin 7 cyuc 2 kcaak 0 tcan 0 dwat 2 m hkucc hmimhtut cyutli tcuphja7n hwicli hrasjlutnyikfrip hpa7tli prikklwis hpa7tmwatli fyikre7t hjonli

drutta Filipendula kamschatica is a PERENNIAL growing to 2 m (6ft) by 2 m (6ft). See above for USDA hardiness. It is hardy to UK zone 3. It is in flower in July. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies, beetles. The plant is self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcaakta Edible Parts: Leaves RootEdible Uses: Young shoots - raw[74]. Leaves - cooked[74]. Root - raw or cooked[74].

nrakta E. Asia - Japan, Siberia. ksacta High grass meadows[74] along streams in the mountains[187].

propta Seed - best sown in the autumn in a cold frame[1]. The seed can also be sown in a cold frame in spring, germinating best at a temperature of 10 - 13°c[200]. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer if they have grown enough. If not, keep them in a cold frame for the winter and plant them out in late spring. Division in autumn or winter[188, 200]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring.

nwatta Requires a humus-rich moist soil in semi-shade[1, 200]. Succeeds in full sun only if the soil is reliably moist throughout the growing season[200]. Dislikes dry or acid soils. Does well in marshy soils. Grows well in heavy clay soils. Plants are hardy to about -25°c[187], when growing in shallow water they even survive if the water is frozen[245]. A good bee plant, the flowers also attract flies. The flowers have a powerful, sweet scent when inhaled from a distance, though there are fishy undertones when inhaled near to[245]. Members of this genus are rarely if ever troubled by browsing deer[233].

3.0.7 Lythrum salicaria

Purple Loosestrife

3 tlin 10 cyuc 2 kcaak 0 tcan 3 dwat 1 m hkucc hmimhtut cyutli tcuphja7nh56s hwicli hrasjlutnyikfrip hpa7tli flatprikklwis hpa7tmwatli xre7t hjonli

drutta Lythrum salicaria is a PERENNIAL growing to 1 m (3ft 3in) by 0.5 m (1ft 8in) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 3. It is in flower from June to August, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy)

and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can grow in water.

kcaakta Edible Parts: Leaves RootEdible Uses: ColouringLeaves - cooked[13, 46, 61, 105]. Rich in calcium[179]. Root - cooked[13]. An edible dye is obtained from the flowers[13].

tcanta Cosmetic Dye Preservative Tannin TeethA decoction of the plant is impregnated into wood, rope etc to prevent it rotting in water[74]. The leaves contain about 12% tannin, the stems 10.5%, the flowers 13.7% and the roots 8.5%[218]. It is probably these tannins that preserve the wood etc[218]. The powdered plant is used cosmetically in face-packs to counteract reddened skin[7].

dwatta Antibiotic Antidiarrhoeal Astringent Dysentery Eczema Hypoglycaemic Styptic VulneraryPurple loosestrife is an astringent herb that is mainly employed as a treatment for diarrhoea and dysentery. It can be safely taken by people of all ages and has been used to help arrest diarrhoea in breast-feeding babies[254]. It can also be used to treat heavy periods and inter-menstrual bleeding[254]. Modern research has shown the whole plant to be antibiotic and to be particularly effective against the micro-organism that causes typhus[254]. The flowering plant is antibiotic, highly astringent, hypoglycaemic, styptic and vulnerary[4, 7, 9, 21, 152, 218, 238]. It is valued as an intestinal disinfectant, especially in cases of enteritis[7], an infusion is used internally in the treatment of diarrhoea, internal bleeding, excessive menstruation etc[9, 222, 238]. The flowering plant is harvested in the summer and can be used fresh or dried[9]. Externally, the plant is used as a cleansing and healing wash for wounds, sores, impetigo, eczema, excess vaginal discharge, vaginal itching etc[7, 222, 238, 254]. The powdered plant is used as a haemostatic in cases of severe nosebleeds[7]. The stems are regarded as gum stimulators and are given to children to chew in order to strengthen weak or bleeding gums[7].

nrakta Europe, including Britain, south to N. Africa east to western and northern Asia. N. America.. ksacta Reed swamps at the margins of lakes and slow-flowing rivers, fens and marshes, avoiding acid soils[17].

propta Seed - sow in the autumn or the spring in a cold frame. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer. If you have sufficient seed it could be worthwhile trying a sowing in situ in the autumn or the spring. Division in March or October[188]. Larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring. Basal cuttings in the spring[238]. Harvest the shoots with plenty of underground stem when they are about 8 - 10cm above the ground. Pot them up into individual pots and keep them in light shade in a cold frame or greenhouse until they are rooting well. Plant them out in the summer.

nwatta Landscape Uses:Container, Massing, Rock garden, Seashore, Specimen. Succeeds in ordinary garden soil, especially if it is damp[1]. Prefers a neutral to alkaline soil[238]. Grows well in heavy clay soils. Grows well in marshy soils[24] and succeeds in shallow water at the edges of ponds[56]. Succeeds in full sun or partial shade[188]. A very hardy plant, tolerating temperatures down to at least -25°c[187]. This species can be very invasive and has been declared a noxious weed in some countries. Since being introduced in N. America it has invaded native marshlands, forming large areas of dense stands and crowding out many native species[274]. A very ornamental plant[1]. A good bee and butterfly plant[24]. Plants usually self-sow when well sited[200]. Special Features:Invasive, Naturalizing, Attracts butterflies, Suitable for cut flowers. In garden design, as well as the above-ground architecture of a plant, root structure considerations help in choosing plants that work together for their optimal soil requirements including nutrients and water. The root pattern is rhizomatous with underground stems sending roots and shoots along their length [2-1].

3.0.8 Caltha leptosepala

Western Marsh Marigold, Howell's marsh marigold, Sulphur marsh marigold

3 tlin 7 cyuc 2 kcaak 0 tcan 1 dwat 0.3 m hkucc

hmmihtut cyutli hja7nhs6s hwicli hrasjlutnykfrp hpa7tli flat-prik hpa7tmwatli xre7t hjonli

drutta *Caltha leptosepala* is a PERENNIAL growing to 0.3 m (1ft) by 0.3 m (1ft in). See above for USDA hardiness. It is hardy to UK zone 3. It is in flower from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, beetles, flies. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It can grow in semi-shade (light woodland) or no shade. It prefers wet soil and can grow in water.

pcista The whole plant, but especially the older portions, contains the toxic glycoside protoanemonin - this is destroyed by heat[172]. The sap can irritate sensitive skin[172].

kcakta Edible Parts: Flowers Leaves RootEdible Uses: Root - it must be well cooked[172]. Some caution is advised, see the notes above on toxicity. Flower buds - raw, cooked or pickled and used as a caper substitute[85, 105, 172, 183]. Some caution is advised, see the notes above on toxicity. Young leaves, before the flowers emerge are eaten raw or cooked[85, 172, 183]. Some caution is advised, see the notes above on toxicity. Older leaves, before the plant flowers, can be eaten if well cooked[85]. Some caution is advised, see the notes above on toxicity.

dwatta Antispasmodic Expectorant Poultice WartsThe whole plant is antispasmodic and expectorant. It has been used to remove warts[172]. A poultice of the chewed roots has been applied to inflamed wounds[257].

nrakta Western N. America - Alaska to Oregon. ksacta Open, wet, subalpine and alpine marshes, wet seepages and marshy meadows at elevations of 750 - 3900 metres[60, 85, 270].

propta Seed - best sown as soon as it is ripe in a cold frame in late summer[200]. Stand the pots in 2 - 3cm of water to keep the soil wet. The seed usually germinates in 1 - 3 months at 15°C[138]. When they are large enough to handle, prick the seedlings out into individual pots and grow them on in a tray of water in a cold frame until they are at least 15cm tall. Plant them out into their permanent positions in the summer. Division in early spring or autumn[200]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the summer or following spring.

nwatta A plant of the waterside, it prefers growing in a sunny position in wet soils or shallow water[1, 111, 233], though it will tolerate drier conditions if there is shade from the summer sun[233]. It requires a deep rich slightly acidic soil[111, 200]. Grows well in heavy clay soils. Plants are hardy to about -20c[187]. A greedy plant, inhibiting the growth of nearby plants, especially legumes[54].

3.0.9 *Caltha leptosepala howellii*

3 tlin 7 cyuc 2 kcak 0 tcan 0 dwat 0.3 m hkuc
hmmihtut cyutli hja7nhs6s hwicli hrasjlutnykfrp hpa7tli flat-prik hpa7tmwatli xre7t hjonli

drutta *Caltha leptosepala howellii* is a PERENNIAL growing to 0.3 m (1ft) by 0.3 m (1ft in). See above for USDA hardiness. It is hardy to UK zone 3. It is in flower from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, beetles, flies. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It can grow in semi-shade (light woodland) or no shade. It prefers wet soil and can grow in water.

pcista The whole plant, but especially the older portions, contains the toxic glycoside protoanemonin - this is destroyed by heat[172]. The sap can irritate sensitive skin[172].

kcakta Edible Parts: Flowers Leaves RootEdible Uses: Root - cooked. A survival food, used when all else fails[172]. Some caution is advised, see the notes above on toxicity. Flower buds - cooked. Added to stews etc or pickled as capers[172]. Some caution is advised, see the notes above on toxicity. Young leaves - cooked. Best before the plant flowers[172]. Some caution is advised, see the notes above on toxicity.

nrakta Western N. America - Alaska to California. ksacta Wet alpine and sub-alpine places[60]. Bogs and wet ground by streams, 1500 - 3000 metres[187].

propta Seed - best sown as soon as it is ripe in a cold frame in late summer[200]. Stand the pots in 2 - 3cm of water to keep the soil wet. The seed usually germinates in 1 - 3 months at 15°C[138]. When they are large enough to handle, prick the seedlings out into individual pots and grow them on in a tray of water in a cold frame until they are at least 15cm tall. Plant them out into their permanent positions in the summer. Division in early spring or autumn[200]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the summer or following spring.

nwatta A plant of boggy soils and shallow pond margins[1], it requires a deep rich slightly acidic soil[200]. It grows well in heavy clay soils. Plants are hardy to about -20°C[187]. A greedy plant, inhibiting the growth of nearby plants, especially legumes[54]. There is some disagreement over the correct name for this species, with some authorities calling it *C. howellii*.

3.0.10 *Caltha palustris*

Marsh Marigold, Yellow marsh marigold
3 tlin 7 cyuc 2 kcak 0 tcan 2 dwat 0.3 m hkuc
hmmihtut cyutli hja7nhs6s hwicli hrasjlutnykfrp hpa7tli flat-prikhwis hpa7tmwatli xre7t hjonli

drutta *Caltha palustris* is a PERENNIAL growing to 0.3 m (1ft) by 0.3 m (1ft in). See above for USDA hardiness. It is hardy to UK zone 3. It is in flower from March to July. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, beetles, flies. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers wet soil and can grow in water.

pcista The whole plant, but especially the older portions, contains the toxic glycoside protoanemonin - this is destroyed by heat[172]. The sap can irritate sensitive skin[172].

kcakta Edible Parts: Flowers Leaves RootEdible Uses: Root - must be well cooked[183]. The raw root should not be eaten[K]. Some caution is advised, see the notes above on toxicity. Flower buds - raw, cooked or pickled and used as a caper substitute[142, 183]. Eating the raw flower buds can lead to intoxication[222]. Some caution is advised, see the notes above on toxicity. Young leaves - raw or cooked[2, 46, 55, 62, 161]. The leaves are harvested in the spring as the plant is coming into flower and is used like spinach after cooking in two or more changes of water[2, 183]. Eating the raw leaves can lead to intoxication[222]. Some caution is advised, see the notes above on toxicity. Older leaves, before the plant flowers, can be eaten if they are well cooked[172]. Some caution is advised, see the notes above on toxicity.

tcanta DyeA yellow dye is obtained from the flowers[[100, 115], a saffron substitute[142]. It is used as a dye when mixed with alum, though it is not very permanent[4]. Plants can be grown for ground cover when planted about 45cm apart each way[208].

dwatta Anodyne Antirheumatic Antispasmodic Diaphoretic Diuretic Expectorant Irritant Rubefacient WartsEvery part of this plant is strongly irritant and so it should be used with caution[4]. The whole plant is anodyne, antispasmodic, diaphoretic, diuretic, expectorant and rubefacient[21, 172]. It has been used to remove warts and is also used in the treatment of fits and anaemia[4, 21, 172]. The root is antirheumatic, diaphoretic, emetic and expectorant[218, 222]. A decoction is used in the treatment of colds[257]. A poultice of the boiled and mashed roots has been applied to sores[257]. A tea made from the leaves is diuretic and laxative[222, 257]. All parts of the plant can irritate or blister the skin or mucous membranes[222].

nrakta Much of the northern Temperate zone in Europe, including Britain, N. America and Asia. ksacta Wet areas in marshes, fens, ditches and wet alder woods[17, 19]. Rare on very base poor peat[17].

propta Seed - best sown as soon as it is ripe in a cold frame in late summer[200]. Stand the pots in 2 - 3cm of water to keep the soil wet. The seed usually germinates in 1 - 3 months at 15°C[138]. When they are large enough to handle, prick the seedlings out into individual pots and grow them on in a tray of water in a cold frame until they are at least

another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them out in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. This species has an excellent potential as a fruit crop in Britain. The fruit is about the size of a cherry, it is very freely borne and the best forms are of excellent dessert quality[K]. The tree is very easily grown and is little troubled by pests or diseases. It also requires very little attention, once the trees are established virtually the only work needed is to harvest the fruit each year[K]. Grafted specimens can produce fruit in their third year[K]. A very ornamental plant[74], it grows well in Britain flowering and fruiting well at Kew and Wisley[K]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Hybridizes freely with other members of this genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

5.2.5 *Crataegus azarolus*

Azarole
5 tlin 9 cyuc 4 kcak 0 tcan 2 dwat 10 m hkuc
hmmihtut cyutli hgattcuphja7n hwicli hrasjlutnykfrp hpa7tli flatprikwlisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus azarolus* is a deciduous Tree growing to 10 m (32ft 10in) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower in June. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked in pies, preserves etc[1, 2, 3, 11, 34, 183]. The fruit can be used fresh or dried for later use. A pleasant acid taste[89]. In warm temperate areas the fruit develops more fruit sugars and has a fragrant sugary pulp with a slightly acid flavour[183]. It can be eaten out of hand. In cooler zones, however, the fruit does not develop so well and is best cooked or used in preserves[183]. The fruit is very variable in size and colour, it is up to 25mm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine

as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta S. Europe to W. Asia. ksacta Dry hillsides and mountains in woods and hedges[50].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. The azarole has long been cultivated for its edible fruit in S. Europe, though it is now going out of favour[3, 11]. There are some named varieties[46]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Hawthorns in general hybridize freely with other members of the genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

5.2.6 *Crataegus pinnatifida major*

Chinese Haw
5 tlin 9 cyuc 4 kcak 2 tcan 3 dwat m hkuc
hmmihtut cyutli hgattcuphja7n hwicli hrasjlutnykfrp hpa7tli flatprikwlisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus pinnatifida major* is a deciduous Tree growing to 7.5 m (24ft 7in). See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender. It is in flower in May. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[3, 105, 177]. A pleasant flavour, it is sold in local markets in China[178]. The fruit can be used in making pies, preserves, etc. and can also be dried for later use. A good size, it is up to 25mm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. This is a tree with an excellent potential as a fruit crop in Britain. The fruit is of very good quality and is freely borne, whilst the tree is of very easy cultivation and rarely troubled by pests or diseases[K]. A tree at the Hillier Arboretum in September 1993 was about 3 metres tall and 1.5 metres wide. The growth looked somewhat weak and the tree was leaning due to wind rock but it was carrying a heavy crop of fruit[K]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. A very ornamental plant[1], it is very closely related to *C. mollis*, and considered to be part of that species by many botanists[229]. Hybridizes freely with other members of this genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

5.2.3 *Crataegus ellwangeriana*

Scarlet Hawthorn
5 tlin 7 cyuc 5 kcaak 2 tcan 2 dwat 6 m hkuac
hmimhtut cyutli hgattcuphja7n hwicli hrasjutnykifrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus ellwangeriana* is a deciduous Tree growing to 6 m (19ft) by 6 m (19ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. Juicy and acid[82]. An excellent dessert fruit, it has an acid yet sweet flavour, is juicy but with a mealy texture, it makes very pleasant eating[K]. The fruit can also be used in making pies, preserves etc, and can be dried for later use. A reasonable size, it is about 15mm in diameter[200]. A specimen at Kew has fruits up to 25mm in diameter[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America. ksacta Not known in the wild, it is possibly of hybrid origin.

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to

pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. This species is possibly of hybrid origin, involving *C. pedicellata*, though some botanists view it as no more than part of *C. pedicellata*[82]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A very ornamental plant[11].

5.2.4 *Crataegus schraderana*

Blue hawthorn
5 tlin 9 cyuc 5 kcaak 0 tcan 2 dwat 6 m hkuac
hmimhtut cyutli hgattcuphja7n hwicli hrasjutnykifrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus schraderana* is a deciduous Tree growing to 6 m (19ft 6in). It is not frost tender. It is in flower in June, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Fruit - raw or cooked[74]. A reasonable size, about 15 - 20mm in diameter[K]. This is one of the nicest tasting fruits of the genus I have tried to date. When fully ripe it is juicy with an extremely pleasant flavour and almost literally melts in the mouth[K]. I would far rather eat this fruit than a strawberry[K]. It ripens in late September and hangs on the tree in good condition for at least 4 weeks[K]. The fruit is so soft that it is best eaten fresh from the tree[K]. The fruit can also be used in making pies, preserves etc, and can be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Although no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta S. Europe - Greece. ksacta Mountainous areas[50].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take

15cm tall. Plant them out into their permanent positions in the summer. Division in early spring or autumn[200]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the summer or following spring.

nwatta A plant of the waterside, it prefers growing in a sunny position in wet soils or shallow water up to 15cm deep[1, 111, 233], though it will tolerate drier conditions if there is shade from the summer sun[233]. Another report says that it grows best in partial shade[17]. Requires a deep rich slightly alkaline soil[31, 111]. Grows well in heavy clay soils. Prefers a chalky soil[31]. Tolerates a pH in the range 4.8 to 7.5. A very ornamental[1] and polymorphic plant[17], there are some named varieties[233]. Plants often self-sow when well sited[208]. A good bee plant[24]. A greedy plant, inhibiting the growth of nearby plants, especially legumes[54]. This species is probably the most primitive flower in the British flora[17].

3.0.11 *Salix alaxensis*

Feltleaf Willow
3 tlin 7 cyuc 1 kcaak 2 tcan 2 dwat 10 m hkuac
htut cyutli tcuphja7n hwicli hrasjutnykifrip hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta *Salix alaxensis* is a deciduous Tree growing to 10 m (32ft 10in). See above for USDA hardiness. It is hardy to UK zone 3 and is not frost tender. It is in flower in May, and the seeds ripen in June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil.

kcakta Edible Parts: Inner bark Leaves NectarEdible Uses: Inner bark - raw or cooked. A sweet flavour[172], it has been eaten as a winter titbit[257]. The taste is somewhat like watermelon or cucumber[257]. The bark has been used as a survival food[229]. It can be dried, ground into a powder and used as a thickening in soups or can be added to cereal flour for use in making bread etc. Young tender leaves and shoots - raw or cooked[257]. The shoots are peeled and eaten in spring[172]. A source of vitamin C[257]. The flowers have been sucked by children for the sweet nectar[257].

tcanta In northern parts of this plant's range, its wood is often the sole source of firewood[229]. Dynamic accumulator.

dwatta Anodyne Febrifuge Vitamin CThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta North-western N. America - Alaska to British Columbia, east to Hudson Bay. ksacta Moist tundra and gravels[172, 204]. Prefers rocky soils along the sides of rivers and lakes[226]. Grows up to and beyond the tree-line, when it becomes a small recumbent low-spreading shrub[226].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta We have very little information on this species and do not know if it will be hardy in Britain, though judging by its native range it should succeed outdoors in most parts of this country. It is an aggregate species[172] and ranges in habit from a small tree right down to a dwarf shrub growing along the ground in exposed sites[229]. The following notes are based on the general needs of the genus. Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Plants

in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

3.0.12 *Athyrium niponicum*

Painted Fern, Japanese Silver Painted Fern
3 tlin 8 cyuc 1 kcaak 0 tcan 0 dwat 0.3 m hkuac
pcimhmim cyutli tcuphja7n hwicli hrasjutnykifrip hpa7tli frusflatprikwih hpa7tmwatli hjonli

drutta *Athyrium niponicum* is a FERN growing to 0.3 m (1ft) by 0.3 m (1ft in) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 4. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils, prefers well-drained soil and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very acid soils. It can grow in full shade (deep woodland) or semi-shade (light woodland). It prefers moist or wet soil.

pcista Although we have found no reports of toxicity for this species, a number of ferns contain carcinogens so some caution is advisable[200]. Many ferns also contain thiaminase, an enzyme that robs the body of its vitamin B complex. In small quantities this enzyme will do no harm to people eating an adequate diet that is rich in vitamin B, though large quantities can cause severe health problems. The enzyme is destroyed by heat or thorough drying, so cooking the plant will remove the thiaminase[172].

kcakta Edible Parts: LeavesEdible Uses: Very young fronds (croziers) - boiled[177]. Some caution is advised. See the notes above on toxicity.

nrakta E. Asia - China, Japan, Korea. ksacta Shaded places in lowland all over Japan[58].

propta Spores - surface sow in a pot of sterile compost in a shady part of the greenhouse and keep moist, this is most easily done by putting the pot in a plastic bag. Pot up small clumps of the plants when they are large enough to handle and keep them moist until they are established. Plant out in late spring of the following year. Division in spring as plants come into growth. Larger divisions can be planted straight into their permanent positions whilst smaller clumps are best potted up and kept in a cold frame until they are growing away well.

nwatta Landscape Uses:Alpine garden, Border, Container, Massing, Rock garden, Woodland garden. An easily grown plant, it is calcifuge and prefers an acid soil with a pH from 4.5 to 6.5, but it tolerates alkaline soils if plenty of leaf mould is added[200]. Grows well in heavy clay soils. Prefers a moist sheltered site with moderately high atmospheric humidity[200]. Succeeds in a semi-shaded bog-garden or in damp woodland, also in garden borders in full or part shade[200]. Members of this genus are rarely if ever troubled by browsing deer[233]. Special Features: Attractive foliage.

4 4 tlin

4.1 bwon

4.1.1 *Alnus viridis crispa*

American Green Alder
tfo2n 4 tlin 8 cyuc 1 kcaak 3 tcan 2 dwat 3 m hkuac
hmimhtut cyutli tcuphja7n hwicli hysijlutnykifrip hpa7tli flatprikwih hpa7tmwatli hjonli

drutta *Alnus viridis crispa* is a deciduous Shrub growing to 3 m (9ft) by 3 m (9ft). See above for USDA hardiness. It is hardy to UK zone 4. It is in flower from April to May. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. It can fix Nitrogen. Suitable for: medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcakta Edible Parts: FlowersEdible Uses: Catkins - raw or cooked. A bitter taste[172].

tcanta Dye TeethAn orange-red to brown dye can be obtained from the bark[257]. Nitrogen fixer.

dwatta Abortifacient Appetizer Astringent Emetic Em-nagogue Haemostatic Poulitice Skin Stomachic Tonic-The bark is astringent, emetic, haemostatic, stomachic and tonic[172]. The bark was burnt as an inhalant in the treatment of rheumatism[257]. The ashes were also used as a tooth cleaner[257]. A decoction of the inner bark has been used as a carminative to reduce gas in the stomach and as a febrifuge[257]. A decoction of the plant has been used in a steam treatment to bring about menstruation - it has been used as an abortifacient[257]. A poultice of the leaves has been used to treat infected wounds or sores[257]. The poultice was left in place over the wound until the leaves stuck to it and was then pulled off, removing the 'poison' with it[257]. An infusion of the plant tops was given to children with poor appetites[257].

nrakta Eastern N. America - Labrador to Alaska and Newfoundland and southwards. ksacta Rocky shores, slopes and mountains[43]. Singly or in thickets along streams, lakeshores, coasts, and bog or muskeg margins, or on sandy or gravelly slopes or flats, from sea level to 2000 metres[270].

propta Seed - best sown in a cold frame as soon as it is ripe and only just covered[200]. Spring sown seed should also germinate successfully so long as it is not covered[200, K]. The seed should germinate in the spring as the weather warms up. When large enough to handle, prick the seedlings out into individual pots. If growth is sufficient, it is possible to plant them out into their permanent positions in the summer, otherwise keep them in pots outdoors and plant them out in the spring. If you have sufficient quantity of seed, it can be sown thinly in an outdoor seed bed in the spring[78]. The seedlings can either be planted out into their permanent positions in the autumn/winter, or they can be allowed to grow on in the seed bed for a further season before planting them. Cuttings of mature wood, taken as soon as the leaves fall in autumn, outdoors in sandy soil.

nwatta Prefers a heavy soil and a damp situation[1, 11]. Grows well in heavy clay soils. A useful plant for cold damp places[11]. Tolerates lime and very infertile sites[11, 200]. This species has a symbiotic relationship with certain soil micro-organisms, these form nodules on the roots of the plants and fix atmospheric nitrogen. Some of this nitrogen is utilized by the growing plant but some can also be used by other plants growing nearby[200]. In garden design, as well as the above-ground architecture of a plant, root structure considerations help in choosing plants that work together for their optimal soil requirements including nutrients and water. The root pattern is branching: a heart root, dividing from the crown into several primary roots going down and out [2-1].

4.2 mran

4.2.1 Crataegus pedicellata

Scarlet Haw, Scarlet hawthorn

4 tlin 8 cyuc 5 kcaak 2 tcan 2 dwat 7 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus pedicellata is a deciduous Shrub growing to 7 m (23ft) by 7 m (23ft). See above for USDA hardness. It is hardy to UK zone 5 and is not frost tender. It is in flower from April to May, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[K]. A very variable species, especially in its fruit which ranges considerable in size and quality[K]. Up to 10mm long and pear shaped, the flesh is sweet but dry and mealy[200]. Another report says that the fruit is up to 20mm in diameter[235] and we have often seen fruits this size, or even a bit larger[K]. We find the fruits of the best forms to have an excellent flavour and juicy texture when fully ripe,

but with a hint of bitterness before then[K]. They are one of the later members of this genus to ripen their fruit[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Northeastern N. America - Connecticut to Ontario, Illinois and Pennsylvania. Naturalized in Britain. ksacta Self-sown in hedges and rough ground in south-eastern England.

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Plants are hardy to at least -18°C[202]. A very variable species, it has been divided up into several different species by some botanists[82]. The sub-species *C. pedicellata gloriosa*. Sarg. has larger fruits than the type[11]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A user in Poland says that *Crataegus pedicellata* exists in the wild in Eastern Poland and can support winter temperatures at least until -30C.

4.2.2 Crataegus missouriensis

4 tlin 8 cyuc 5 kcaak 0 tcan 2 dwat m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwli hpa7tmwatli xre7t hjonli

drutta Crataegus missouriensis is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardness. It is hardy to UK zone 5 and is not frost tender. The seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for:

5.2 mran

5.2.1 Crataegus pennsylvanica

5 tlin 9 cyuc 5 kcaak 0 tcan 2 dwat 9 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus pennsylvanica is a deciduous Tree growing to 9 m (29ft 6in). See above for USDA hardness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Fruit - raw or cooked. A very good size fruit, up to 25mm in diameter, with a lovely semi-sweet flavour even when semi-ripe, it makes a delicious dessert[K]. It is also sometimes made into jellies[82] and can be dried for later use. The fruit is borne in clusters of 4 - 12 and is very easily harvested[82, K]. There are up to five fairly large seeds in the centre of each fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Although no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Delaware to Pennsylvania and W. Virginia. ksacta Open woods and thickets in moist fertile soils[43]. Meadows in low, moist soils[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. This species has a good potential as a fruit crop in Britain. It regularly produces heavy crops of high quality, the tree is very easily grown and is little troubled by pests or diseases. It also requires very little attention, once the trees are established virtually the only work needed is to harvest the fruit each year[K]. A tree about 5 metres tall and wide at Kew fruits regularly

and very heavily[K]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

5.2.2 Crataegus arnoldiana

Arnold Hawthorn

5 tlin 9 cyuc 5 kcaak 0 tcan 2 dwat 7 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus arnoldiana is a deciduous Tree growing to 7 m (23ft). See above for USDA hardness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[105, 177]. Sub-acid[82]. A delicious flavour, it is sweet with a soft juicy flesh and makes an excellent dessert fruit[K]. It can also be cooked and used in pies, preserves etc and can be dried for later use. The fruit ripens in early September in southern Britain[K]. The fruit is about 2cm in diameter[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta North-eastern N. America - Massachusetts and Connecticut. ksacta Wooded banks[43]. Thickets on a dry bank[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in

nwatta Landscape Uses:Border, Container, Foundation, Massing, Rock garden, Specimen, Woodland garden. Easily grown in ordinary garden soil[1]. It does well in a moist heavy soil in full sun or partial shade[1, 17, 31, 90] and is happy in water-logged conditions[90]. The dormant bulbs are fairly hardy and will withstand soil temperatures down to at least -5°C[214]. Plants can be naturalized in damp rough grass[200]. Flowers are produced within 4 - 5 years from seed. The seedpods are swollen and filled with air so that they can be dispersed by floating in water[90]. Members of this genus are rarely if ever troubled by browsing deer or rabbits[233]. Special Features:Attractive foliage, Naturalizing, Suitable for cut flowers.

4.2.55 Lysimachia nummularia

Creeping Jenny, Moneywort, Creeping Charlie
4 tin 8 cyuc 1 kcak 0 tcan 2 dwat 0.2 m hkuc
hmmihtut cyutli tcuphja7nhs6s hwicli hrsajlutnykfrp
hpa7tli flatpriklwis hpa7tmwatli fyfxre7t hjonli

drutta Lysimachia nummularia is an evergreen Perennial growing to 0.2 m (0ft 8in) by 0.6 m (2ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 4. It is in leaf all year, in flower from June to August, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies. The plant is self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can grow in water.

kcakta Edible Parts: Edible Uses: TeaA tea is made from the leaves and flowers[46, 61, 177, 183].

tcanta Can be used as a ground cover plant in moist soils but requires weeding for the first year or so. Plants should be spaced about 60cm apart each way and form a carpet, rooting as they spread[208].

dwatta Antidiarrhoeal Antiscorbatic Astringent Diuretic VulneraryThe whole herb, used fresh or dried, is antiscorbatic, mildly astringent, diuretic and vulnerary[4, 9, 46]. For drying, the herb is harvested in June[4]. Creeping Jenny is a very good wound-herb, the fresh leaves being bruised and applied externally to the affected area[4]. An infusion is used to treat internal bleeding and diarrhoea[9].

nrakta Europe, including Britain, from Sweden south and east to Spain, N. Greece and the Caucasus. ksacta Damp situations in pastures and by the sides of streams, avoiding acid soils[4, 9].

propta Seed - sow spring or autumn in a cold frame[200]. This species rarely produces seed in Britain[17]. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer. Division in spring or autumn[200]. Larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring.

nwatta Landscape Uses:Container, Ground cover, Rock garden, Woodland garden. An easily grown plant, succeeding in a moist loamy soil[1]. Grows well in heavy clay soils. Grows well in marshy soil and in shallow water in pond margins[24, 56]. Prefers a shady position[111] but also succeeds in full sun[188]. Plants are hardy to at least -25°C[187]. Most species in this genus seem to be immune to the predations of rabbits[233]. A very ornamental plant[1]. Special Features:Attractive foliage, Invasive, Naturalizing, Fragrant flowers, Inconspicuous flowers or blooms.

5 5 tlin

5.1 bwon

5.1.1 Alnus tenuifolia

Mountain Alder, Thinleaf alder
tfo2n 5 tlin 7 cyuc 1 kcak 0 tcan 2 dwat 9 m hkuc
hmmihtut cyutli tcuphja7n hwicli hysijlutnykfrp hpa7tli flat-
priklwis hpa7tmwatli hjonli

drutta Alnus tenuifolia is a deciduous Tree growing to 9 m (29ft 6in) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 2. It is in flower in March, and the seeds ripen in October. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. It can fix Nitrogen. Suitable for: medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista The freshly harvested inner bark is emetic but is alright once it has been dried[172].

kcakta Edible Parts: FlowersEdible Uses: Catkins - raw or cooked. A bitter taste[172].

tcanta Pioneer Soil stabilization Tannin WoodThis is an excellent pioneer species for re-establishing woodlands on disused farmland, difficult sites etc[226]. Its fast rate of growth means that it quickly provides sheltered conditions to allow more permanent woodland trees to become established. In addition, bacteria on the roots fix atmospheric nitrogen - whilst this enables the tree to grow well in quite poor soils it also makes some of this nitrogen available to other plants growing nearby. Alder trees also have a heavy leaf canopy and when the leaves fall in the autumn they help to build up the humus content of the soil. Alder seedlings do not compete well in shady woodland conditions and so this species gradually dies out as the other trees become established[K]. The tree has an extensive root system and can be planted to control banks from erosion[226]. The bark and the strobils are a source of tannin[82]. A dark dye is obtained from the bark[226]. The colour can range from orange through red to brown[257]. Wood - soft, straight-grained, very durable in water[82]. It is of no commercial value, though it is used locally as a fuel[229].

dwatta Anodyne Astringent Emetic Febrifuge Haemostatic Stomachic TonicThe bark is astringent, emetic, haemostatic, stomachic and tonic[172]. The bark also contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226]. The outer bark is astringent and is applied as a poultice to bleeding wounds, it also reduces swellings[226].

nrakta Western N. America - Alaska to California and New Mexico. ksacta Moist soils by swamps, streams, ponds and lakes in foothills to well up in the mountains[60, 229].

propta Seed - best sown in a cold frame as soon as it is ripe and only just covered[200]. Spring sown seed should also germinate successfully so long as it is not covered[200, K]. The seed should germinate in the spring as the weather warms up. When large enough to handle, prick the seedlings out into individual pots. If growth is sufficient, it is possible to plant them out into their permanent positions in the summer, otherwise keep them in pots outdoors and plant them out in the spring. If you have sufficient quantity of seed, it can be sown thinly in an outdoor seed bed in the spring[78]. The seedlings can either be planted out into their permanent positions in the autumn/winter, or they can be allowed to grow on in the seed bed for a further season before planting them. Cuttings of mature wood, taken as soon as the leaves fall in autumn, outdoors in sandy soil.

nwatta Landscape Uses: Erosion control. Prefers a heavy soil and a damp situation[1, 11]. Grows well in heavy clay soils[11]. Tolerates very infertile sites[200]. A fast-growing but short-lived tree[229]. There is some confusion over the correct name of this tree with one authority citing the European species *A. incana* as the correct name[60]. Another report says that this species is closely related to *A. incana*, but distinct[229]. Some modern works treat it as a subspecies (*Alnus incana tenuifolia*). This species has a symbiotic relationship with certain soil micro-organisms, these form nodules on the roots of the plants and fix atmospheric nitrogen. Some of this nitrogen is utilized by the growing plant but some can also be used by other plants growing nearby[200]. Special Features:Not North American native, Wetlands plant, Inconspicuous flowers or blooms. In garden design, as well as the above-ground architecture of a plant, root structure considerations help in choosing plants that work together for their optimal soil requirements including nutrients and water. The root pattern is branching: a heart root, dividing from the crown into several primary roots going down and out [2-1].

light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. The size of a large cherry, the flesh is sweet, soft and fairly juicy[K]. The fruit is only about 10mm thick according to another report[235]. The fruit ripens in mid to late September in southern Britain[K], it makes a delicious dessert fruit and can also be used in making pies, preserves etc[K]. The fruit can be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotoxic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta South-eastern N. America - Virginia to North Carolina and Missouri. ksacta Rocky bluffs and river banks[235].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. This species has a good potential as a fruit crop in Britain. It regularly produces heavy crops of high quality, the tree is very easily grown and is little troubled by pests or diseases. It also requires very little attention, once the trees are established virtually the only work needed is to harvest the fruit each year[K]. A plant growing at Hillier Arboretum was about 3 metres tall (with a trunk 2 metres long) and 3 metres wide, it was fruiting well in 1993 and 1994[K]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.3 Quercus robur

Pedunculate Oak, English oak
4 tin 8 cyuc 4 kcak 5 tcan 3 dwat 30 m hkuc
hmmihtut cyutli hgattcuphja7n hwicli jlutnykfrp hpa7tli flat-
priklwis hpa7tmwatli hjonli

drutta Quercus robur is a deciduous Tree growing to 30 m (98ft) by 30 m (98ft) at a slow rate. See above for USDA hardiness. It is hardy to UK zone 4. It is in flower from April to May, and the seeds ripen from September to October. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. It is noted for attracting wildlife. Suitable for: medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure.

pcista Possible digestive complaints. May delay absorption of alkaloids and other alkaline drugs [301].

kcakta Edible Parts: SeedEdible Uses: Coffee Gum-Seed - cooked[2, 5, 8, 13]. Nourishing but indigestible[4]. Chopped and roasted, the seed is used as an almond substitute[8]. It can be dried, ground into a powder and used as a thickening in stews etc or mixed with cereals for making bread[183]. The seed contains bitter tannins, these can be leached out by thoroughly washing the seed in running water though many minerals will also be lost[63]. Either the whole seed can be used or the seed can be dried and ground it into a powder. It can take several days or even weeks to properly leach whole seeds, one method was to wrap them in a cloth bag and place them in a stream. Leaching the powder is quicker. A simple taste test can tell when the tannin has been leached. The traditional method of preparing the seed was to bury it in boggy ground overwinter. The germinating seed was dug up in the spring when it would have lost most of its astringency. The roasted seed is a coffee substitute[21, 61]. An edible gum is obtained from the bark[177]. Another report says that an edible manna is obtained from the plant and that it is used instead of butter in cooking[183]. This report probably refers to the gum[K].

tcanta Basketry Charcoal Compost Fuel Gum Ink Repellent Tannin WoodA mulch of the leaves repels slugs, grubs etc, though fresh leaves should not be used as these can inhibit plant growth[20, 201]. The bark is an ingredient of 'Quick Return' herbal compost activator[32]. This is a dried and powdered mixture of several herbs that can be added to a compost heap in order to speed up bacterial activity and thus shorten the time needed to make the compost[K]. The bark is very rich in calcium[18]. Oak galls are excrescences that are sometimes produced in great numbers on the tree and are caused by the activity of the larvae of different insects. The insects live inside these galls, obtaining their nutrient therein. When the insect pupates and leaves, the gall can be used as a rich source of tannin, that can also be used as a dyestuff[4]. A black dye and an excellent long-lasting ink is made from the oak galls, mixed with salts of iron[4, 7, 66]. The colour is not very durable[4]. When mixed with alum, the dye is brown and with salts of tin it is yellow[4]. Trees can be coppiced to provide material for basket making, fuel, construction etc[23]. The wood is a source of tar, quaiacol, acetic acid, creosote and tannin[123]. Tannin is extracted commercially from the bark and is also found in the leaves[223]. On a 10% moisture basis, the bark contains 11.6% tannin and the wood 9.2%[223]. The bark strips easily from the wood in April and May[4]. A purplish dye is obtained from an infusion of the bark with a small quantity of copperas[4]. It is not bright, but is said to be durable[4]. Wood - hard, tough, durable even under water - highly valued for furniture, construction etc[4, 13, 61, 66]. It is also a good fuel[6] and charcoal[61].

dwatta Antiseptic Astringent Bach Decongestant Dysentery Haemostatic TonicThe oak tree has a long history of medicinal use. It is anti-inflammatory, antiseptic, astringent, decongestant, haemostatic and tonic[4, 7, 9, 13, 21, 165]. The bark is the part of the plant that is most commonly used[4], though other parts such as the galls, seeds and seed cups are also sometimes used[7]. A decoction of the bark is useful in the treatment of chronic diarrhoea, dysentery, intermittent fevers, haemorrhages etc[4]. Externally, it is used to bathe wounds, skin eruptions, sweaty feet, piles etc[9]. It

is also used as a vaginal douche for genital inflammations and discharge, and also as a wash for throat and mouth infections[9]. The bark is harvested from branches 5 - 12 years old, and is dried for later use[9]. Any galls produced on the tree are strongly astringent and can be used in the treatment of haemorrhages, chronic diarrhoea, dysentery etc[4]. The plant is used in Bach flower remedies - the keywords for prescribing it are 'Despondency', 'Despair, but never ceasing effort'[209]. A homeopathic remedy is made from the bark. It is used in the treatment of disorders of the spleen and gall bladder[9]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve *Quercus robur* Pedunculate Oak for coughs/bronchitis, diarrhoea, inflammation of mouth and pharynx, inflammation of the skin (see [302] for critics of commission E).

nrakta Europe, including Britain, from Scandinavia south and east to Spain, the Urals and Crimea. ksacta Often the dominant woodland tree, especially on clay soils and in the eastern half of Britain, but avoiding acid peat and shallow limestone soils[17].

propta Seed - it quickly loses viability if it is allowed to dry out. It can be stored moist and cool overwinter but is best sown as soon as it is ripe in an outdoor seed bed, though it must be protected from mice, squirrels etc. Small quantities of seed can be sown in deep pots in a cold frame. Plants produce a deep taproot and need to be planted out into their permanent positions as soon as possible, in fact seed sown in situ will produce the best trees[11]. Trees should not be left in a nursery bed for more than 2 growing seasons without being moved or they will transplant very badly.

nwatta Prefers a good deep fertile loam which can be on the stiff side[11]. Young plants tolerate reasonable levels of side shade[200]. Succeeds in heavy clay soils[13] and in wet soils so long as the ground is not water-logged for long periods[186]. Dislikes dry or shallow soils but is otherwise drought tolerant once it is established[186]. Tolerant of exposed sites though it dislikes salt-laden winds[186]. The oak is a very important timber tree in Britain, it is also a very important food plant for the caterpillars of many species of butterfly[30], there are 284 insect species associated with this tree[24]. It has often been coppiced or pollarded for its wood in the past[23], though this should not be done too frequently[186], about once every 50 years is the average. The tree flowers on new growth produced in spring, the seed ripening in its first year[200, 229]. Older trees have a thick corky bark and this can protect them from forest fires, young trees will often regenerate from the base if cut down or killed back by a fire[186]. Intolerant of root disturbance, trees should be planted in their permanent positions whilst young[11]. Hybridizes freely with other members of the genus[200]. Immune to attacks by the tortix moth[1]. This species is notably resistant to honey fungus[88, 200].

4.2.4 Quercus bicolor

Swamp White Oak

4 tlin 8 cyuc 4 kcak 3 tcan 2 dwat 25 m hkuc hmimhtut cyutli tcuphja7n hwicli jltunyikfrip hpa7tli flat-priklwis hpa7tmwatli hjonli

drutta *Quercus bicolor* is a deciduous Tree growing to 25 m (82ft 0in) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower in May, and the seeds ripen in October. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. Suitable for: medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure.

kcakta Edible Parts: SeedEdible Uses: CoffeeSeed - raw or cooked[105, 161, 226]. A rather sweet flavour[159, 235]. The seed is quite large, about 2 - 3cm long and 15 - 20mm wide[82, 200], and unlike most other oaks, is attached to the tree by a long stem[183]. It matures in its first year[235]. The seed can be dried, ground into a powder and used as a thickening in stews etc or mixed with cereals for making bread[183]. The seed from some trees can contain bitter tannins, these can be leached out by thoroughly washing the seed in running water though many minerals will also be

lost. Either the whole seed can be used or the seed can be dried and ground it into a powder. It can take several days or even weeks to properly leach whole seeds, one method was to wrap them in a cloth bag and place them in a stream. Leaching the powder is quicker. A simple taste test can tell when the tannin has been leached. The traditional method of preparing the seed was to bury it in boggy ground overwinter. The germinating seed was dug up in the spring when it would have lost most of its astringency. Roasted seed is a coffee substitute.

tcanta Repellent Rust Tannin WoodThe leaves of most species in this genus are more or less rich in tannins. A mulch of the leaves can be placed around vulnerable plants in order to repel slugs, snails, grubs etc. Fresh leaves should be used with caution, since these can utilize some of the nitrogen in the soil and this inhibit plant growth[20 , K]. Oak galls are excrescences that are sometimes produced in great numbers on the tree and are caused by the activity of the larvae of different insects. The insects live inside these galls, obtaining their nutrient therein. When the insect pupates and leaves, the gall can be used as a rich source of tannin, that can also be used as a dyestuff[4]. The bark has been boiled with hemlock (*Tsuga* sp.?) and soft maple bark (*Acer* spp), and the liquid used to remove rust. The mixture was also believed to prevent rust[257]. The wood is close-grained, strong, hard, tough, heavy[46 , 82 , 171 , 229]. It weighs 48lb per cubic foot[235]. Trees do not self-prune and dead branches remain on the trees for many years. This does mean that the wood is usually quite knotty[226]. The wood machines well, though it can check and warp if not dried properly. It is of some importance commercially. And is used for construction, cabinet making, furniture, interior finishes, veneers, fence posts etc[6 , 82 , 171 , 229 , 338 , 1050]. The wood makes a good fuel.

dwatta Astringent DysenteryAny galls produced on the tree are strongly astringent and can be used in the treatment of haemorrhages, chronic diarrhoea, dysentery etc[4].

nrakta Eastern N. America - Quebec to Minnesota, Georgia and Arkansas. ksacta Bottomlands, stream margins and swamps[43]. Tolerant of poorly drained sites, it is frequently found in heavy mucky soils[229].

propta Seed - it quickly loses viability if it is allowed to dry out. It can be stored moist and cool overwinter but is best sown as soon as it is ripe in an outdoor seed bed, though it must be protected from mice, squirrels etc. Small quantities of seed can be sown in deep pots in a cold frame. Plants produce a deep taproot and need to be planted out into their permanent positions as soon as possible, in fact seed sown in situ will produce the best trees[11]. Trees should not be left in a nursery bed for more than 2 growing seasons without being moved or they will transplant very badly.

nwatta Landscape Uses:Pest tolerant, Specimen, Street tree. Prefers a good deep fertile loam which can be on the stiff side[1, 11]. Young plants tolerate reasonable levels of side shade[200]. Tolerates moderate exposure, surviving well but being somewhat stunted[200]. Prefers warmer summers than are usually experienced in Britain, trees often grow poorly in this country and fail to properly ripen their wood resulting in frost damage overwinter[200]. A relatively fast-growing tree in the wild, living 300 - 350 years[229], though it is relatively shallow-rooted and is considerably more susceptible to forest fires than most other oaks[226]. This species is one of the best of the white oaks for growing in this country[1]. However, it is said that the seed is rarely formed in Britain[105]. The tree flowers on new growth produced in spring, the seed ripening in its first year[200, 229]. Trees do not commence bearing until about 25 - 30 years old, large crops being produced every 3 - 5 years[229]. Intolerant of root disturbance, trees should be planted in their permanent positions whilst young[11]. The seed of this species is attached to the tree by unusually long (by oak tree standards) stems[183]. Hybridizes freely with other members of the genus[200]. Plants in this genus are notably resistant to honey fungus[200]. Special Features: Attracts birds, North American native, Attracts butterflies, Inconspicuous flowers or blooms. The plant is heat tolerant in zones 8 through 1. (Plant Hardiness Zones show how well plants withstand cold winter temperatures. Plant Heat Zones show when plants would start suffering from the heat. The Plant Heat Zone map is based on the number of "heat days" experienced in a given area where the temperature climbs to over 86 degrees F (30°C). At this temperature, many plants begin to

are about 8 - 10cm above the ground. Pot them up into individual pots and keep them in light shade in a cold frame or greenhouse until they are rooting well. Plant them out in the summer.

nwatta An easily grown plant, succeeding in a moist or wet loamy soil in sun or partial shade[187]. Prefers a shady position[111]. Grows well in heavy clay soils. Hardy to at least -25°C[187]. Most species in this genus seem to be immune to the predations of rabbits[233]. A very ornamental plant[1]. The sub-species *L. vulgaris davurica*. (Ledeb.)Kunth. is the form used for food in China and Japan[177].

4.2.52 Juncus effusus

Soft Rush, Common rush, Lamp rush, Pacific rush

4 tlin 8 cyuc 1 kcak 0 tcan 2 dwat 1.5 m hkuc hmimhtut cyutli tcuphja7nhs6s hwicli hrasjlutnyikfrip hpa7tli flatprik hpa7tmwatli xre7t hjonli

drutta *Juncus effusus* is a PERENNIAL growing to 1.5 m (5ft) by 0.5 m (1ft 8in). See above for USDA hardiness. It is hardy to UK zone 4. It is in flower from June to August. The species is hermaphrodite (has both male and female organs) and is pollinated by Wind. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can grow in water. The plant can tolerate strong winds but not maritime exposure.

pcista Possibly toxic to mammals[76].

kcakta Edible Parts: LeavesEdible Uses: Young shoots - raw[118, 257]. Some caution is advised, see the notes above on toxicity.

tcanta Basketry Lighting Paper Strewing String Thatching WeavingStems are used in basket making, thatching, weaving mats etc[23, 46, 66, 99, 115, 171]. The stems can also be dried then twisted or braided into ropes for tying or binding[257]. Stems can be peeled (except for a small spine which is left to keep them upright) and soaked in oil then used as a candle[6, 46, 61, 115]. A fibre obtained from the stems is used for making paper[189]. The stems are harvested in late summer or autumn, they are split and cut into usable pieces and then soaked for 24 hours in clear water. They are then cooked for 2 hours with lye and beaten in a blender. The fibres make an off-white paper[189]. When mixed with mulberry fibres they can be used for making stencil paper[189]. The whole plant was formerly used as a strewing herb[1, 66, 115].

dwatta Antiphlogistic Depurative Diuretic Febrifuge Lenitive Lithontrpic Pectoral Sedative UrinaryThe pith of the stem is antiphlogistic, depurative, discutient, diuretic, febrifuge, lenitive, lithontrpic, pectoral and sedative[147, 176, 178, 218, 240]. It is used in the treatment of sore throats, jaundice, oedema, acute urinary tract infection and morbid crying of babies[176].

nrakta Throughout the northern temperate zone, including Britain, east and south Africa, Australasia. ksacta Wet pastures, bogs, damp woods etc, usually on acid soils[17].

propta Seed - surface sow in pots in a cold frame in early spring and keep the compost moist. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer if they have grown sufficiently, otherwise in late spring of the following year. Division in spring. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring.

nwatta Easily grown in a moist soil, bog garden or shallow water[1, 200]. Prefers a heavy soil in sun or light shade[200].

4.2.53 Iris sibirica

Siberian Iris

4 tlin 9 cyuc 1 kcak 0 tcan 1 dwat 0.6 m hkuc hmimhtut cyutli tcuphja7n hwicli hrasjlutnyikfrip hpa7tli flat-priklwis hpa7tmwatli fyfjxre7t hjonli

drutta *Iris sibirica* is a PERENNIAL growing to 0.6 m (2ft) by 1 m (3ft 3in) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 6 and is not frost tender.

It is in flower in June, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Insects. The plant is self-fertile. Suitable for: light (sandy) and medium (loamy) soils, prefers well-drained soil and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista Many plants in this genus are thought to be poisonous if ingested, so caution is advised[65]. The roots are especially likely to be toxic[238]. Plants can cause skin irritations and allergies in some people[238].

kcakta Edible Parts: RootEdible Uses: Root - an edible starch is said to be extracted from it[2, 105, 177]. The root is not eaten in China[178]. Some caution is advised since there are reports that the plant might be poisonous. Perhaps the extracted starch is edible.

tcanta InsecticideAn insecticide is obtained from the plant[178]. (from the root?)

dwatta ExpectorantExpectorant[178].

nrakta C. Europe to Russia. ksacta Damp woodland[50], pastures and heaths[100].

propta Seed - best sown as soon as it is ripe in a cold frame. Stored seed should be sown as early in the year as possible in a cold frame. Prick out the seedlings into individual pots when they are large enough to handle and grow them on in the greenhouse or cold frame for their first year. Plant out into their permanent positions in late spring or early summer. Division, best done in August/September after flowering but can also be done in April. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring.

nwatta Landscape Uses:Border, Container, Foundation, Massing, Specimen, Woodland garden. Prefers a humus-rich soil[79], succeeding in a moist border or by water[42, 79]. Grows well in heavy clay soils. The plants are very cold tolerant, but they can be damaged when dormant if the soil is too moist[79]. A very ornamental plant, there are many named varieties[187]. Plant the roots out very shallowly[79]. Members of this genus are rarely if ever troubled by browsing deer or rabbits[233]. Special Features: Attractive foliage, Not North American native, Naturalizing, All or parts of this plant are poisonous, Wetlands plant, Suitable for cut flowers.

4.2.54 Leucojum aestivum

Summer Snowflake

4 tlin 9 cyuc 1 kcak 0 tcan 0 dwat 0.5 m hkuc hmimhtut cyutli tcuphja7n hwicli hrasjlutnyikfrip hpa7tli prikklwis hpa7tmwatli xre7t hjonli

drutta *Leucojum aestivum* is a BULB growing to 0.5 m (1ft 8in) by 0.1 m (0ft 4in) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower in May, and the seeds ripen in July. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcakta Edible Parts: RootEdible Uses: Bulb - cooked[105]. I have some reservations about this report, though no records of the plant being poisonous have been found so far.

nrakta Europe, including Britain, from France south and east to Spain, Greece, the Crimea and Iran. ksacta Marshes, wet meadows and willow thickets, avoiding acid soils[17].

propta The seed is best sown as soon as it is ripe in a well-drained soil in a cold frame. Stored seed requires 2 - 3 months cold stratification, it should then germinate in 2 - 4 weeks at 10°C[134]. Sow the seed thinly so that the seedlings can be allowed to grow on undisturbed in the pots for their first year of growth. Give them an occasional weak liquid feed to ensure that they do not become nutrient deficient. Pot up the small bulbs when dormant, planting 2 - 3 bulbs in each pot. Grow them on for another 2 - 3 years before planting them out. Division of offsets in September/October. It is best done as soon as the foliage ripens[1]. Scooping the bulbs.

is cultivated for its branches which are used in basket making[11], there are some named varieties[131]. Plants are coppiced annually for this purpose[186] A very important food plant for the caterpillars of many butterfly species[30] and a good bee plant, providing an early source of nectar and pollen[11]. Plants in this genus are notably susceptible to honey fungus[200]. Plants should be put into their permanent positions as soon as possible[11]. Dioecious. Male and female plants must be grown if seed is required.

4.2.49 Salix commutata

undergreen willow
4 tin 8 cyuc 1 kcaak 3 tcan 2 dwat 3 m hkuc
hmimhtut cyutli tcuphja7n hwicli hrasjlutnykfrkip hpa7tli flatprik hpa7tmwtli nyanhjen hjonli

drutta Salix commutata is a deciduous Shrub growing to 3 m (9ft 10in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower in May. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - raw or cooked. They are not very palatable[172].

tcanta Soil reclamationA pioneer plant for difficult sites[200]. Dynamic accumulator.

dwatta Anodyne FebrifugeThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta Western N. America - Alaska to California. ksacta Wet places at moderate to rather high elevations[204].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

4.2.50 Salix brachycarpa

shortfruit willow
4 tin 8 cyuc 1 kcaak 3 tcan 2 dwat 1.4 m hkuc
htut cyutli tcuphja7n hwicli hrasjlutnykfrkip hpa7tli flatprik hpa7tmwtli nyanhjen hjonli

drutta Salix brachycarpa is a deciduous Shrub growing to 1.4 m (4ft 7in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate maritime exposure.

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread

etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - not very palatable[172].

tcanta ShelterbeltA salt tolerant coastal plant, it can be grown as part of a shelterbelt[200]. Dynamic accumulator.

dwatta Anodyne FebrifugeThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta N. America - Canada to Oregon. S. Greenland. ksacta Found in a wide range of habitats, including saline, calcareous and serpentine, from lowland to alpine elevations[204].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on chalk[200]. Very tolerant of maritime exposure[200]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

4.2.51 Lysimachia vulgaris

Yellow Loosestripe, Garden yellow loosestripe
4 tin 8 cyuc 1 kcaak 0 tcan 2 dwat 1.2 m hkuc
hmimhtut cyutli tcuphja7nhs6s hwicli hrasjlutnykfrkip hpa7tli flatprikwis hpa7tmwtli fyfxire7t hjonli

drutta Lysimachia vulgaris is a PERENNIAL growing to 1.2 m (4ft). See above for USDA hardiness. It is hardy to UK zone 5. It is in flower from April to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Bees, flies. The plant is self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can grow in water.

kcakta Edible Parts: LeavesEdible Uses: Young leaves[105, 177]. No more details are given.

tcanta DyeA yellow dye is obtained from the flowers[13, 100]. A brown dye is obtained from the rhizomes[13]. The growing plant repels gnats and flies, it has been burnt in houses in order to remove these insects[4].

dwatta Antidiarrhoeal Astringent Demulcent Dysentery ExpectorantAn astringent herb, yellow loosestripe is principally used to treat gastro-intestinal conditions such as diarrhoea and dysentery, to stop internal and external bleeding and to cleanse wounds[254]. The herb is astringent, demulcent and expectorant[4, 61]. It is harvested when in flower in July and dried for later use[4]. The plant can be used internally or externally and is useful in checking bleeding of the mouth, nose and wounds, restraining profuse haemorrhages of any kind and in the treatment of diarrhoea[4]. It makes a serviceable mouthwash for treating sore gums and mouth ulcers[254].

nrakta Most areas of Europe and Asia, including Britain, but excluding the extreme north and south. ksacta Marshes, streams and in shallow water in reed swamps[187]. Shady places near water, avoiding acid soils[4, 17].

propta Seed - sow spring or autumn in a cold frame. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer. Division in spring or autumn[200]. Very easy, larger clumps can be replanted direct into their permanent positions, though it is best to pot up smaller clumps and grow them on in a cold frame until they are rooting well. Plant them out in the spring. Basal cuttings, March to April in a cold frame. Harvest the shoots with plenty of underground stem when they

suffer physiological damage. Heat Zones range from 1 (no heat days) to 12 (210 or more heat days). For example Heat Zone. 11-1 indicates that the plant is heat tolerant in zones 11 through 1.) For polyculture design as well as the above-ground architecture (form - tree, shrub etc. and size shown above) information on the habit and root pattern is also useful and given here if available. The plant growth habit is a standard with a non-suckering single trunk [1-2].

4.2.5 Crataegus holmesiana

Holmes' hawthorn
4 tin 8 cyuc 4 kcaak 0 tcan 2 dwat 10 m hkuc
hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrkip hpa7tli flatprikwislip hpa7tmwtli xre7t hjonli

drutta Crataegus holmesiana is a deciduous Tree growing to 10 m (32ft) by 8 m (26ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. An excellent dessert fruit, it has an acid yet sweet flavour, is juicy but with a mealy texture, it makes very pleasant eating[K]. Some forms have fruits with a disagreeable flavour[82]. The fruit can also be used in making pies, preserves etc, and can be dried for later use. The fruit is about 10mm in diameter[200]. A specimen at Cambridge Botanical garden has fruits up to 20mm in diameter[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiac HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Ontario to Maine, south to New York and Pennsylvania. ksacta Rich moist hillsides and the borders of streams and swamps[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on

a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hawthorns in general hybridize freely with other members of the genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A very ornamental plant[11].

4.2.6 Crataegus douglasii

Black Hawthorn
4 tin 8 cyuc 4 kcaak 0 tcan 2 dwat 9 m hkuc
hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrkip hpa7tli flatprikwislip hpa7tmwtli xre7t hjonli

drutta Crataegus douglasii is a deciduous Tree growing to 9 m (29ft 6in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[2, 3, 95, 99, 101]. A very pleasant flavour with a sweet and juicy succulent flesh[46, 82, K], it makes an excellent dessert fruit and can be eaten in quantity[K]. The fruit can also be used for making pies, preserves etc, and can be dried for later use[183]. The fruit is about 8mm in diameter and is borne in small clusters[200]. The fruits I have eaten have been considerably larger than this[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Needles WoodThe spines on the branches are used as needles for lancing boils, removing splinters etc[99]. Wood - close-grained, heavy, hard and tough. Used for tool handles etc[82, 99, 101].

dwatta Antirheumatic Astringent Cardiotonic Dysentery Hypotensive Poulitce StomachicAn infusion of the shoots has been used to treat diarrhoea in children and sores in babies mouths[257]. A poultice of the chewed leaves has been applied to swellings[257]. An infusion of the bark has been used in the treatment of diarrhoea and dysentery[257]. An infusion of the sapwood, bark and roots has been used as a stomach medicine[257]. The thorns have been used as a treatment for arthritis[257].The point of the thorn was used to pierce an area affected by arthritic pain. The other end of the thorn was ignited and burned down to the point buried into the skin. This treatment was very painful but it was said that after a scab had formed and disappeared, the arthritic pain had also disappeared[257]. The thorns have been used as probes for boils and ulcers[257]. Although no other specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Western N. America - British Columbia to Michigan, south to California. ksacta Open woods, banks of mountain streams and on rocky banks[43, 82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it

should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.7 Crataegus mollis

Red Haw, Downy hawthorn
4 tlin 8 cyuc 4 kcaak 2 tcan 2 dwat 6 m hkuc
hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrp hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus mollis is a deciduous Tree growing to 9 m (29ft) by 12 m (39ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from May to June, and the seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: TeaFruit - raw or cooked[22, 46, 62, 101, 183]. Sub-acid, dry and mealy[82, 227]. The fruit is also used for making jellies and preserves[61, 95, 183]. Fairly large, it is about 20 - 25mm in diameter[200], with a thick flesh[229]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K]. A tea-like beverage can be made from the twigs[257].

tcanta Hedge HedgeThe plant is tolerant of being trimmed and will even resprout if cut back into very old wood. It can be used as a hedge[50]. Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern and Central N. America - Ontario to Alabama, west to Oklahoma. ksacta Open woods, usually in alluvial or fertile soils[43]. Frequently found in limestone soils[229].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta Landscape Uses:Espalier, Pollard, Specimen, Street tree. A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. A very hardy plant, tolerating temperatures down to at least -18°C[202]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A very ornamental plant[11], the branches are almost thornless[229]. A rather variable species, it has been separated into several different species by some botanists[82]. This species is much confused with *C. chrysoarpa*[11] and with *C. coccinea*[50]. It is very closely related to *C. submollis*, differing in having 20 stamens to a flower (as compared to 10 in *C. submollis*)[202]. Special Features: North American native, Attractive flowers or blooms. The plant is heat tolerant in zones 6 through 1. (Plant Hardiness Zones show how well plants withstand cold winter temperatures. Plant Heat Zones show when plants would start suffering from the heat. The Plant Heat Zone map is based on the number of "heat days" experienced in a given area where the temperature climbs to over 86 degrees F (30°C). At this temperature, many plants begin to suffer physiological damage. Heat Zones range from 1 (no heat days) to 12 (210 or more heat days). For example Heat Zone. 11-1 indicates that the plant is heat tolerant in zones 11 through 1. For polyculture design as well as the above-ground architecture (form - tree, shrub etc. and size shown above) information on the habit and root pattern is also useful and given here if available. The plant growth habit is a standard with a non-suckering single trunk [1-2]. The root pattern is a tap root similar to a carrot going directly down [1-2].

4.2.8 Crataegus acclivis

4 tlin 8 cyuc 4 kcaak 0 tcan 2 dwat 8 m hkuc
hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnykfrp hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus acclivis is a deciduous Tree growing to 8 m (26ft) by 7 m (23ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from April to May, and the seeds ripen in September. The species

4.2.47 Salix viminalis

Osier. Basket Willow
4 tlin 8 cyuc 1 kcaak 3 tcan 2 dwat 6 m hkuc
htut cyutli tcuphja7n hwicli hrasjlutnykfrp hpa7tli flatprikwisslip hpa7tmwatli nyanhjen hjonli

drutta Salix viminalis is a deciduous Tree growing to 6 m (19ft) by 4 m (13ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower from April to May, and the seeds ripen in June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required). and is pollinated by Bees. The plant is not self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - cooked. Not very palatable[172].

tcanta Basketry Fodder Soil stabilization TanninThe stems are very flexible and are used in basket making[11, 23, 46]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. The annual yield can be around 12 tonnes per hectare, 40% of which is class 1[74]. The bark contains about 10% tannin[223]. Often planted along the banks of rivers and lakes to prevent soil erosion[11].

dwatta Anodyne Antirheumatic FebrifugeAntirheumatic, febrifuge[13]. The fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta Europe, incl Britain, from Scandinavia south and east to France through temperate Asia to Japan. ksacta By rivers and streams, also on deep moist alluvial soils, avoiding very acid soils[186].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Fodder: Bank Industrial Crop: Biomass Industrial Crop: Tannin Management: Coppice Other Systems: SRC Regional CropSucceeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Dislikes heavy shade and dry soils[186]. Rarely thrives on chalk[200]. Tolerates atmospheric pollution[186]. A fast growing tree, it is very wind resistant[K]. Widely cultivated for basket making, there are many named varieties[11, 17]. Trees are usually coppiced annually for this purpose, they do not respond so well to pollarding. It should be cut back almost to ground level each year when being grown for basket making since this encourages the production of long flexible shoots[11]. In very good conditions plants can put on 4 metres or more of new growth in a year when treated in this way. A very important food source for the caterpillars of many butterfly species[30], it is also a valuable early pollen source for bees[11, 186]. The plants are rich in insect life[186]. Best planted into its permanent position as soon as possible[11]. The root system is rather aggressive and can cause problems with drains[200]. Plants are best not grown within 10 metres of buildings. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

4.2.48 Salix purpurea

Purple Osier. purpleosier willow
4 tlin 8 cyuc 1 kcaak 4 tcan 3 dwat 5 m hkuc

htut cyutli kyucutpghja7n hwicli hrasjlutnykfrp hpa7tli flatprikwihssom hpa7tmwatli nyanhjen hjonli

drutta Salix purpurea is a deciduous Tree growing to 5 m (16ft) by 5 m (16ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from March to April, and the seeds ripen in May. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required). and is pollinated by Bees. The plant is not self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in saline soils. It cannot grow in the shade. It prefers dry moist or wet soil.

pcista Gastrointestinal bleeding and kidney damage possible. Avoid concurrent administration with other aspirin-like drugs. Avoid during pregnancy. Drug interactions associated with salicylates applicable [301].

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - raw or cooked. They are not very palatable[172].

tcanta Basketry Fodder Hedge Hedge Repellent Soil reclamation Soil stabilization TanninThe stems are very tough and flexible and are used in basket making[11, 13, 23, 115]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. The bark is much disliked by rabbits, so a closely woven fence of this plant can be used as a protective barrier[115]. The bark contains about 10% tannin[223]. Plants can be grown as a hedge[29], the var. 'Gracilis' is suitable for a small hedge on damp sites[182, 200]. It can be kept dense by annual clipping[11]. The plant has an extensive root system and is used in soil reclamation and stabilization projects along estuaries[199]. Dynamic accumulator.

dwatta Anodyne Antinflammatory Antiperiodic Antispastic Astringent Diaphoretic Diuretic Dysentery Febrifuge Hypnotic Sedative TonicThe bark is anodyne, anti-inflammatory, antiperiodic, antispetic, astringent, diaphoretic, diuretic, febrifuge, hypnotic, sedative and tonic[4, 9, 21]. It is a very rich source of salicin, which is used in making aspirin[199]. The bark of this species is used interchangeably with *S. alba*. It is taken internally in the treatment of rheumatism, arthritis, gout, inflammatory stages of auto-immune diseases, diarrhoea, dysentery, feverish illnesses, neuralgia and headache[238]. The bark is removed during the summer and dried for later use[238]. The leaves are used internally in the treatment of minor feverish illnesses and colic[238], cancerous sores and chronic dysentery[218]. The leaves can be harvested throughout the growing season and are used fresh or dried[238]. The twigs are used in the treatment of cancer, dysentery and ulcers[218]. The bark of the stem and roots is anodyne and stypitic[218]. It is used in the treatment of rheumatism[218]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve *Salix* / Willow for diseases accompanied by fever, rheumatic ailments, headaches (see [302] for critics of commission E).

nrakta Europe, including Britain, from Belgium south and east to N. Africa, temperate Asia to Japan. ksacta Wet places in lowland areas[13, 17], preferring neutral or alkaline soils[186].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Agroforestry Services: Windbreak Fodder: Bank Industrial Crop: Biomass Management: Coppice Other Systems: SRC Regional CropSucceeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Plants prefer an alkaline or neutral soil, rarely doing well in acid conditions[186]. Said to prefer a sandy soil[23, 199], plants are tolerant of dryish soils[11]. Plants are tolerant of salt water[199]. A very ornamental plant[1, 11]

astringent[240]. A gum and the juice of the trees are used to increase visual powers[240].

nrakta Europe, including Britain, from Norway south and east to Spain, temperate Asia and Syria. ksacta Woods, scrub and hedges, usually on basic soils, to 840 metres[17].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Cuttings of this species do not root well[200]. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Cuttings of this species do not root well[200].

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1], but prefers a damp, heavy soil in a sunny position. Grows in drier soils than any other British species of Salix[186]. Rarely thrives on chalk[200]. Plants are found most frequently on basic soils in the wild[17]. Tolerates atmospheric pollution and exposed positions, including maritime exposure[75, 186]. A fast growing tree, it establishes well[75]. The tree has an untidy habit[75]. A light demanding tree, it becomes tall and drawn when grown in woodland, though it grows well along the sunnier edges[186]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Trees are very tolerant of cutting, they coppice well[186]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required. Special Features:Not North American native, Wetlands plant, Attractive flowers or blooms.

4.2.45 Salix acutifolia

Sharp-Leaf Willow

4 tin 8 cyuc 1 kcaak 3 tcan 2 dwat 10 m hkuc

htut cyutli tucpja7n hwicli hrasljutnykfrp hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta Salix acutifolia is a deciduous Shrub growing to 10 m (32ft 10in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required). and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate maritime exposure.

kcakta Edible Parts: Inner bark LeavesEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and added to cereal flours for use in making bread etc. A famine food, it is only used when all else fails[172]. Young shoots - cooked. They are not very palatable[172].

tcanta Basketry Shelterbelt Soil stabilizationStems are very flexible and are used in basket making[46, 61]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. Trees can be planted in shelter-belts for protection against the wind[166]. The extensive root system of this plant is good for binding sandy soils[11]. Dynamic accumulator.

dwatta Anodyne FebrifugeThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta N. Europe to E. Asia. ksacta Not known

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1, 11], but prefers a damp, heavy soil in a sunny position[200]. Rarely thrives on

chalk[200]. Very wind-resistant, tolerating maritime exposure[166]. Hybridizes freely with other members of this genus[200]. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. Closely related to S. daphnoides[200] and considered to be a part of that species by some authorities[11, 17]. Some named forms have been developed for their ornamental value[182]. There are also named forms cultivated for basket making[46, 61]. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

4.2.46 Salix daphnoides

Violet Willow, Daphne willow

4 tin 8 cyuc 1 kcaak 3 tcan 2 dwat 10 m hkuc

htut cyutli tucpja7n hwicli hrasljutnykfrp hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta Salix daphnoides is a deciduous Tree growing to 10 m (32ft) by 8 m (26ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from February to March, and the seeds ripen from May to June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required). and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate maritime exposure.

kcakta Edible Parts: Flowers Inner bark Leaves RootEdible Uses: Inner bark - raw or cooked[172, 183]. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - raw or cooked. They are not very palatable[172]. Rich in vitamin C[183]. Young catkin-bearing shoots - raw or cooked[183]. The growing tips of underground rhizomes are peeled and eaten[183].

tcanta Basketry Shelterbelt Soil stabilization WoodThe stems are very flexible and are used in basket making[1, 61, 74, 131, 158] or for making wattle and daub walls[158]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. Trees are fast growing and tolerate maritime exposure so can be grown as a shelterbelt[166]. Plants have extensive root systems and are often used to stabilize sand dunes[11, 200]. Wood - soft, even grained, smooth. Used for construction, pales, tubs etc[146, 158]. Dynamic accumulator.

dwatta Anodyne Febrifuge Vitamin CThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226].

nrakta Europe to W. Asia as far as the Himalayas. Scarcely naturalized in Britain[17]. ksacta Moist soils.

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in most soils, including wet, ill-drained or intermittently flooded soils[1], but prefers a deep damp, heavy soil in a sunny position[11, 200]. Rarely thrives on chalk[200]. Succeeds in poor strong soils[131]. Succeeds in maritime exposure[166]. The root system is rather aggressive and can cause problems with drains[200]. It is best not to plant this species within 10 metres of buildings. Hybridizes freely with other members of this genus. Although the flowers are produced in catkins early in the year, they are pollinated by bees and other insects rather than by the wind[11]. A very ornamental and fast growing plant[1, 74]. There are some named varieties. Plants in this genus are notably susceptible to honey fungus[200]. Dioecious. Male and female plants must be grown if seed is required.

is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[K]. The fruit ripens in early September, it is about 18mm long with a pleasantly sweet, juicy flesh and makes an excellent dessert fruit[K]. The fruit contains up to 5 seeds in the centre, these usually stick together and so the effect is like eating a cherry with its single large seed[K].

dwatta Cardiac HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta North-eastern N. America - New York to the borders of southern Canada. ksacta Banks of streams and steep gorges[235].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Plants are hardy to at least -18°C[202]. We have very little specific information on this plant, and it is regarded as no more than a form of C. pedicellata by most botanists[235]. However, a tree seen at Kew in early September 1997 had a good crop of almost ripe fruit. This fruit was more elongated than C. pedicellata and was also ripe about 4 weeks before that species[K]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.9 Crataegus submollis

Quebec Hawthorn

4 tin 8 cyuc 4 kcaak 0 tcan 2 dwat 7 m hkuc

hmmihtut cyutli hgattcuphja7n hwicli hrasljutnykfrp hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta Crataegus submollis is a deciduous Tree growing to 7 m (23ft) by 7 m (23ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in June, and the seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[105, 161]. Sub-acid, dry and mealy[82]. Our experience has been that the fruit is sweet and somewhat juicy, it has a thick flesh with a nice flavour and makes an acceptable dessert fruit[K]. The fruit can also be used in making pies, preserves etc, and can be dried for later use[257]. It is about 20mm in diameter, is borne in large clusters and is easily harvested[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Hedge HedgePlants are very tolerant of trimming and will soon resprout even if cut back into very old wood. They can be used as a hedge[50]. Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta North-eastern N. America - Quebec to Ontario, Massachusetts and New York. ksacta Wooded hillsides and open fultlets ground[43]. Rich damp hillsides and the borders of woods and roads[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Plants are hardy to about -18°C[202]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are

the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. This species is sometimes confused with *C. coccinea*[50]. It is also very close to *C. mollis*, differing in having 10 stamens to a flower (as compared to 20 in *C. mollis*)[202]. It is considered to be part of *C. mollis* by some botanists[229].

4.2.10 Crataegus succulenta

Fleshy hawthorn
4 tin 8 cyuc 4 ckak 2 tcan 2 dwat 6 m hkuc
hmimhtut cyutli hgatkycuctuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli
drutta *Crataegus succulenta* is a deciduous Shrub growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils, prefers well-drained soil and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

ckaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[62, 101, 105]. Excellent for making jellies[183], it can also be dried for later use. A small fruit with a sweet, juicy and pulpy flesh[82, 183], it is up to 17mm in diameter[235]. Another report says that the fruit is large for a *Crataegus* species, it is about 35mm in diameter and is borne in small clusters[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Nova Scotia to North Carolina, west to Montana, Colorado and Utah. ksacata Thickets, pastures and woodland edges, especially in dry or rocky soils[43].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in

full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. This species is closely related to *C. tomentosa*[11].

4.2.11 Crataegus durobrivensis

Caughuawaga Hawthorn
4 tin 8 cyuc 4 ckak 2 tcan 2 dwat 5 m hkuc
hmimhtut cyutli hgatkycuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus durobrivensis* is a deciduous Shrub growing to 5 m (16ft) by 5 m (16ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate atmospheric pollution.

ckaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. A reasonable size, it is up to 15mm in diameter[200], and is very acceptable for raw eating[K]. It is sweet and fairly juicy when fully ripe with a hint of apple in its flavour[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta North-eastern N. America. ksacata Not known in a truly wild situation, it is possibly a hybrid *C. pruinosa* x *C. suborbiculata*[43].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all

nwatta Easily grown in any moist soil in sun or part shade[187, 200]. Prefers a damp position[133, 200]. Grows well in heavy clay soils[208]. Dislikes very light or dry soils[1]. Grows well in the shade of a north-facing wall[208]. Plants are hardy to about -25°C[187]. A very ornamental plant, there are many named varieties[187]. It can be grown in a wild garden, rock garden or the flower border[1].

4.2.43 Salix fragilis

Crack Willow
4 tin 8 cyuc 1 ckak 3 tcan 3 dwat 15 m hkuc
htut cyutli tucphja7n hwicli hrasjlutnyikfrip hpa7tli flatprik hpa7tmwatli nyanhjen hjonli

drutta *Salix fragilis* is a deciduous Tree growing to 15 m (49ft) by 15 m (49ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from April to May, and the seeds ripen from May to June. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid and neutral soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure.

pcista Gastrointestinal bleeding and kidney damage possible. Avoid concurrent administration with other aspirin-like drugs. Avoid during pregnancy. Drug interactions associated with salicylates applicable [301].

ckaakta Edible Parts: Inner bark LeavesEdible Uses: SweetenerInner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - raw or cooked. They are not very palatable[172]. A saccharine exudation is obtained from the leaves and young branches[2, 105]. Used as a food[183].

tcanta Basketry Charcoal Tannin WoodStems are very flexible and are used in basket making[46, 61]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. The bark contains around 10% tannin[223]. Wood - tough, withstands friction. Used for floors, bases of carts etc[11]. A good quality charcoal is obtained from the wood[61]. Dynamic accumulator.

dwatta Anodyne Astringent Dysentery Febrifuge Poulitic StypticThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne, astringent and febrifuge[9, 226]. The bark of this species is used interchangeably with *S. alba*. It is taken internally in the treatment of rheumatism, arthritis, gout, inflammatory stages of auto-immune diseases, diarrhoea, dysentery, feverish illnesses, neuralgia and headaches[238]. A poultice of the bark has been applied to sores as a styptic and healing agent[257]. The bark is removed during the summer and dried for later use[238]. The leaves are used internally in the treatment of minor feverish illnesses and colic[238]. The leaves can be harvested throughout the growing season and are used fresh or dried[238]. The German Commission E Monographs, a therapeutic guide to herbal medicine, approve *Salix* / Willow for diseases accompanied by fever, rheumatic ailments, headaches (see [302] for critics of commission E).

nrakta Europe, including Britain, from Sweden south and east to Spain, Serbia and Iran. ksacata Streambanks, marshes, fens and wet woods[17].

propta Seed - must be surface sown as soon as it is ripe in late spring. It has a very short viability, perhaps as little as a few days. Cuttings of mature wood of the current year's growth, November to February in a sheltered outdoor bed or planted straight into their permanent position and given a good weed-suppressing mulch. Very easy. Plant into their permanent positions in the autumn. Cuttings of half-ripe wood, June to August in a frame. Very easy.

nwatta Succeeds in moist soils, including wet, ill-drained or intermittently flooded soils,[1] but prefers a deep damp, heavy soil in a sunny position[11, 200]. Rarely thrives on

chalk[200]. Succeeds in an exposed position[186]. Closely allied to *S. alba*, with which it freely hybridizes[11], though it tolerates poorer soils than that species[186]. A very important food plant for the caterpillars of many butterfly species[30] and a good bee plant, providing an early source of nectar and pollen[11]. The cultivar 'Basfordiana' is used for basket making[131]. Best if planted into its permanent position as soon as possible. The root system is rather aggressive and can cause problems with drains[200]. Plants in this genus are notably susceptible to honey fungus[200]. Plants are very amenable to coppicing or pollarding and can be cut back annually if required[186]. Dioecious. Male and female plants must be grown if seed is required.

4.2.44 Salix caprea

Goat Willow, Kilmarnock Willow, Pink Pussy Willow, Pussy Willow

4 tin 9 cyuc 1 ckak 4 tcan 2 dwat 10 m hkuc
hmimhtut cyutli kyuctuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikwlis hpa7tmwatli nyanhjen hjonli

drutta *Salix caprea* is a deciduous Tree growing to 10 m (32ft) by 8 m (26ft) at a fast rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from March to April, and the seeds ripen in May. The species is dioecious (individual flowers are either male or female, but only one sex is to be found on any one plant so both male and female plants must be grown if seed is required), and is pollinated by Bees. The plant is not self-fertile. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil. The plant can tolerate maritime exposure. It can tolerate atmospheric pollution.

ckaakta Edible Parts: Inner bark Leaves MannaEdible Uses: Inner bark - raw or cooked. It can be dried, ground into a powder and then added to cereal flour for use in making bread etc. A very bitter flavour, it is a famine food that is only used when all else fails[172]. Young shoots - raw or cooked. They are not very palatable[172]. The source of an edible manna[183]. No further details.

tcanta Basketry Charcoal Hedge Leather Pioneer Shelterbelt Tannin WoodLandscape Uses: Border, Massing, Pollard, Standard, Specimen. The stems are very flexible and are used in basket making[61]. The plant is usually coppiced annually when grown for basket making, though it is possible to coppice it every two years if thick poles are required as uprights. The bark is tough and flexible, it is used as a substitute for leather[115]. The bark contains around 10% tannin[223]. The plant is fast growing and tolerant of maritime exposure, it can be used as a windbreak hedge and shelterbelt though it is of untidy habit[75]. The seeds are very light and so can travel some distance in the wind. The plant is therefore able to find its way to areas such as cleared woodland where the soil has been disturbed. Seedlings will grow away quickly, even in exposed conditions and the plant will provide good shelter for the establishment of woodland plants. Thus it makes a good pioneer species and, except in wetter and moorland-type soils, will eventually be largely out-competed by the other woodland trees. Its main disadvantage as a pioneer plant is that it has an extensive root system and is quite a greedy plant, thus it will not help as much in enriching the soil for the other woodland plants as other pioneer species such as the alders, *Alnus* species[K]. Some cultivars can be grown as ground cover[208]. 'Pendula' is female whilst 'Kilmarnock' is a male, they should be spaced about 1.5 metres apart each way[208]. Wood - soft, elastic, easily split. Used for baskets, rugs etc[46, 61]. A good quality charcoal is made from the wood[46, 61]. Dynamic accumulator.

dwatta Anodyne Aphrodisiac Astringent Febrifuge Ophthalmic StimulantThe fresh bark of all members of this genus contains salicin[226], which probably decomposes into salicylic acid (closely related to aspirin) in the human body[213]. This is used as an anodyne and febrifuge[226]. A decoction of the leaves is used in the treatment of fevers[240]. A distilled water from the flowers is aphrodisiac, cordial and stimulant[240]. It is used externally in the treatment of headaches and ophthalmia[240]. The ashes of the wood are useful in the treatment of haemoptysis[240]. The stems and the leaves are

to handle, prick the seedlings out into individual pots and grow them on in the greenhouse for at least their first winter. Plant them out into their permanent positions in late spring or early summer, after the last expected frosts. Division is possible at almost any time of the year so long as there is sufficient moisture[233]. It is best carried out in March as growth commences or in early autumn if the soil is not too dry[200]. Hostas can be left undivided for many years and should not be divided any more frequently than once every 3 - 5 years to allow the leaves to reach maturity[200].

nwatta Thrives in most fertile soils if they are rich in humus[200]. Grows well in heavy clay soils and in sandy ones[208]. Very limy soils inhibit growth, but plants can thrive in such a situation if plenty of humus is added[208]. Prefers a pH between 6 and 7[200]. Requires a rich soil that does not dry out readily[1]. A plant of boggy places, it is flood tolerant[187]. Does well in the semi-shade of a woodland[200] also succeeding in deeper shade[187]. In general, the sunnier the position the moister the soil should be[233]. Plants are best not grown under trees in town gardens since the snot washed from the leaves of the trees in wet weather will tend to remain on the hosta[208]. Plants flower better when grown in a sunny position but the foliage is better when the plant is in a shady position[200]. Plants are in general fully hardy in Britain, but young leaves in spring can be destroyed by frost. New leaves are only produced in the spring and very early summer, so any damage at this time has a deep effect on the plant[233]. The rhizome creeps extensively[200]. Members of this genus are rarely if ever troubled by browsing deer or rabbits[233]. Plants are very attractive to slugs and snails, the young shoots in spring are especially at risk[111, K]. A polymorphic species, it hybridizes freely with other members of this genus[200].

4.2.40 Eriophorum angustifolium

Cotton Grass, Tall cottongrass
4 tin 8 cyuc 2 kcaok 0 tcan 1 dwat 0.6 m hkuc
hmimhtut cyutli hja7nhs6s hwicli hrasjlutnykfrp hpa7ti frusflat hpa7tmwatli hjonli

drutta Eriophorum angustifolium is a PERENNIAL growing to 0.6 m (2ft) by 1 m (3ft 3in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower from May to June, and the seeds ripen from July to August. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid soils and can grow in very acid soils. It can grow in semi-shade (light woodland) or no shade. It prefers wet soil and can grow in water.

kcakta Edible Parts: Root StemEdible Uses: Young stem bases - raw or cooked[172]. Usually cooked and eaten with oil[257]. Root - raw or cooked[257]. The blackish covering should be removed[172].

tcanta Paper Stuffing Tinder Weaving WickThe cottony seed hairs are used to make candle wicks[4, 13, 100, 172]. They are also used for stuffing pillows[4, 74, 141], paper making etc and as a tinder[74]. Experiments have been made in using the hairs as a cotton substitute, but they are more brittle than cotton and do not bear twisting so well[4]. The dried leaves and stems have been woven into soft mats or covers[257].

dwatta AstringentThe leaves and roots are considerably astringent and have been used in the past as a treatment for diarrhoea[4]. Some native North American Indian tribes would eat the stems raw in order to restore good health to people in generally poor health[257].

nrakta Arctic and temperate regions of Europe, including Britain, to Siberia and N. America. ksacta Peat bogs, acid meadows and marshes[13].

propta Seed - sow in situ in spring in a moist soil in light shade. Germination usually takes place within 2 - 6 weeks at 15°C[200]. If the seed is in short supply it can be sown in pots in a cold frame. Place the pots in a try of water to keep the compost moist. When large enough to handle, prick the seedlings out into individual pots and plant them out in the summer. Division in spring or autumn. Very easy, the divisions can be replanted direct into their permanent positions.

nwatta Requires boggy conditions or a pond margin and an acid soil[1, 162]. Grows well in heavy clay soils. Quite invasive.

4.2.41 Mimulus lewisii

Great Purple Monkey Flower, Purple monkeyflower
4 tin 8 cyuc 2 kcaok 0 tcan 1 dwat 0.6 m hkuc
hmimhtut cyutli tucphja7nhs6s hwicli hrasjlutnykfrp hpa7ti flatprinklwis hpa7tmwatli hjonli

drutta Mimulus lewisii is a PERENNIAL growing to 0.6 m (2ft) by 0.5 m (1ft 8in). See above for USDA hardiness. It is hardy to UK zone 5. It is in flower from July to August. The flowers are pollinated by Insects. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can grow in water.

kcakta Edible Parts: LeavesEdible Uses: Leaves and stems - raw or cooked[172].

dwatta PoulitcePoulitce[172].

nrakta Western N. America - Alaska to Mexico. ksacta Moist areas, stream edges and wet slopes[172]. In and along streams from moderate to high elevations[60].

propta Seed - sow spring in a greenhouse and only lightly cover the seed[200]. When they are large enough to handle, prick the seedlings out into individual pots and plant them out in the summer. If you have sufficient seed it can also be sown in situ in late spring after the last expected frosts[200]. Division in spring[200]. Very easy, large divisions can be planted out direct into their permanent positions. We have found that it is better to pot up the smaller divisions and grow them on in light shade in a cold frame until they are well established before planting them out in late spring or early summer. Cuttings of softwood, 5cm long in spring or summer[200].

nwatta Prefers rather moist places such as damp borders, margins of streams and boggy spots[1]. Grows well in heavy clay soils. Thrives in full sun so long as the soil does not dry out[200], otherwise it is best grown in partial shade[187]. This species tolerates slightly drier conditions than *M. guttatus*, it prefers a rich soil in light woodland[200]. Hardy to at least -20°C[187]. Another report says that plants are only hardy in the milder parts of Britain[233]. There is some confusion over the naming of this plant, the entry on edibility may refer to *M. luteus*, which is closely related to this species.

4.2.42 Primula denticulata

Drumstick Primula
4 tin 8 cyuc 2 kcaok 0 tcan 0 dwat 0.3 m hkuc
hmimhtut cyutli tucphja7n hwicli hrasjlutnykfrp hpa7ti flatprinklwis hpa7tmwatli xre7t hjonli

drutta Primula denticulata is a PERENNIAL growing to 0.3 m (1ft) by 0.2 m (0ft 8in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Insects. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcakta Edible Parts: FlowersEdible Uses: Flowers - raw[105, 177]. They make a very attractive addition to salads[183].

tcanta Plants can be grown as a ground cover, they are best spaced about 35cm apart each way[208].

nrakta E. Asia - Himalayas. ksacta Shrubberies and open slopes in moist situations, often along drains, 1500 - 4500 metres[51, 145].

propta Seed - best sown as soon as it is ripe in a cold frame[133]. Sow stored seed in early spring in a cold frame[1]. When they are large enough to handle, prick the seedlings out into individual pots and grow them on in the greenhouse for their first winter. Plant them out into their permanent positions in late spring or early summer, after the last expected frosts. Root cuttings[200].

fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. A very hardy plant, tolerating temperatures down to about -18°C[202]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. Probably a natural hybrid, *C. pruinosa* x *C. suborbiculata*[43].

4.2.12 Crataegus punctata

Dotted Hawthorn,
4 tin 8 cyuc 3 kcaok 2 tcan 2 dwat 10 m hkuc
hmimhtut cyutli hgattucphja7n hwicli hrasjlutnykfrp hpa7ti flatprinklwis hpa7tmwatli xre7t hjonli

drutta Crataegus punctata is a deciduous Tree growing to 10 m (32ft 10in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower in June, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[159]. Eaten out of hand or made into jellies, preserves etc[177, 183]. A good size, about 25mm in diameter, and borne in small clusters[200]. Ripening in October, it is quite apple-like in texture and appearance, making a good dessert fruit[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta MiscellanyWood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic Hypotensive MiscellanyA compound decoction of the shoots and bark has been used to stop menstrual flow[257]. Although no other specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Quebec to Georgia, west to Minnesota and Oklahoma. ksacta Open rocky ground, thickets and pasture[43]. Rich hillside[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year,

planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A very ornamental plant[1], it grows very well in Britain, flowering and fruiting abundantly[11]. There are some named varieties selected for their ornamental value[200]. 'Ohio Pioneer' is a thornless form growing about 8 metres tall[200]. The plant is heat tolerant in zones 7 through 5. (Plant Hardiness Zones show how well plants withstand cold winter temperatures. Plant Heat Zones show when plants would start suffering from the heat. The Plant Heat Zone map is based on the number of "heat days" experienced in a given area where the temperature climbs to over 86 degrees F (30°C). At this temperature, many plants begin to suffer physiological damage. Heat Zones range from 1 (no heat days) to 12 (210 or more heat days). For example Heat Zone 11-1 indicates that the plant is heat tolerant in zones 11 through 1.) For polyculture design as well as the above-ground architecture (form - tree, shrub etc. and size shown above) information on the habit and root pattern is also useful and given here if available. The plant growth habit is a standard with a non-suckering single trunk [1-2]. The root pattern is a tap root similar to a carrot going directly down [1-2].

4.2.13 Crataegus opaca

Western Mayhaw
4 tin 8 cyuc 3 kcaok 3 tcan 2 dwat 9 m hkuc
hmimhtut cyutli hgattucphja7n hwicli hrasjlutnykfrp hpa7ti frusflatprinklwis hpa7tmwatli xre7t hjonli

drutta Crataegus opaca is a deciduous Shrub growing to 9 m (29ft 6in) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower in March, and the seeds ripen from May to June. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very acid soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[160]. Juicy and acid with a pleasant flavour[2]. It is up to 2cm in diameter[27]. The fruit is frequently used and much prized in parts of southern N. America where it is often gathered in quantity from the wild. Its acid flavour makes it a favourite for preserves and jellies[82, 183, 227]. The fruit can also be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard and strong, but not large enough for commercial use[227]. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine

as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Southern N. America - North Carolina to Mississippi. ksacta Deep depressions filled with water for most of the year, borders of swamps, low river banks etc[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. It thrives in acid soils[160]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. A very hardy species, tolerating temperatures down to about -30°C[160]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. Occasionally cultivated for its fruit in America, there are some named varieties[183].

4.2.14 *Crataegus pedicellata gloriosa*

Scarlet Haw

4 tin 8 cyuc 3 kcaak 3 tcan 2 dwat 7 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnyikfrp hpa7tli flatprikliwsslip hpa7tmwatli xre7t hjonli

drutta *Crataegus pedicellata gloriosa* is a deciduous Shrub growing to 7 m (23ft) by 7 m (23ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from April to May, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[K]. Up to 2cm long and pear shaped[11]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. There are up to five fairly large seeds in

the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - New York. ksacta Not known

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Plants are hardy to at least -18°C[202]. This sub-species has larger fruits than the type[11]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.15 *Malus coronaria*

Garland Crab, Sweet crab apple

4 tin 8 cyuc 3 kcaak 0 tcan 1 dwat 7 m hkuc hmimhtut cyutli tucphja7n hwicli hrasjlutnyikfrp hpa7tli flatprikliwis hpa7tmwatli xre7t hjonli

drutta *Malus coronaria* is a deciduous Tree growing to 7 m (23ft) by 7 m (23ft). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower from May to June, and the seeds ripen from October to November. The species is hermaphrodite (has both male and female organs) and is pollinated by Insects. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils, prefers well-drained soil and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista All members of this genus contain the toxin hydrogen cyanide in their seeds and possibly also in their leaves, but not in their fruits. Hydrogen cyanide is the substance that gives almonds their characteristic taste but it should

The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Quebec to Ontario. ksacta Calcareous ridges and banks of streams[43].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them out in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.38 *Crataegus uniflora*

Dwarf hawthorn

4 tin 8 cyuc 2 kcaak 0 tcan 2 dwat 3 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnyikfrp hpa7tli flatprikliwsslip hpa7tmwatli xre7t hjonli

drutta *Crataegus uniflora* is a deciduous Shrub growing to 3 m (9ft 10in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[161, 177]. The fruit is about 10mm in diameter and is borne in small clusters[200]. The fruit is usually borne singly, or in groups of up to 3 fruits[235]. It can be up to 16mm in diameter and has a firm flesh[235]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers

of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta South-eastern N. America - New York to Florida, west to Texas and Oklahoma. ksacta Sandy or rocky banks or woods[43].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. This species is often confused with *C. tomentosa*. L.[235]. It hybridizes freely with other members of the genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.39 *Hosta longissima*

Swamp Hosta

4 tin 8 cyuc 2 kcaak 0 tcan 0 dwat 0.2 m hkuc hmimhtut cyutli tucphja7n hwicli hrasjlutnyikfrp hpa7tli flatprikliwis hpa7tmwatli xre7t hjonli

drutta *Hosta longissima* is a PERENNIAL growing to 0.2 m (0ft 8in) by 0.5 m (1ft 8in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower from July to August, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs). Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

kcakta Edible Parts: Leaves StemEdible Uses: Young leaves and leaf stems - cooked[105, 177].

nrakta E. Asia - Japan. ksacta Swampy grassland and other wet places, often exposed to full sun in the spring but then shaded in the summer by the long grasses it grows amongst[251].

propta Seed - sow spring in a lightly shaded position in a greenhouse and only just cover the seed. Germination usually takes place within 1 - 3 months at 10°C. Make sure you keep the compost moist. When they are large enough

Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta Landscape Uses:Border, Espalier, Pollard, Screen, Specimen, Street tree. A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. A very hardy plant, tolerating temperatures down to about -18°C[202]. This species is closely related to *C. crus-galli*[11]. It hybridizes freely with other members of the genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. Special Features:Attracts birds, North American native, Blooms are very showy.

4.2.36 *Crataegus rivularis*

River Hawthorn
4 tin 8 cyuc 2 kcaak 0 tcan 2 dwat 4 m hkuc
hmimhtut cyuti hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwlwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus rivularis* is a deciduous Tree growing to 4 m (13ft 1in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[212]. The flesh is thin, dry and mealy[227]. The fruit can also be dried for later use or for making into pemmican[212]. The fruit is high in sugar but low in fats and protein[212]. The fruit is about 10mm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these are stuck together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it

to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Western N. America - Wyoming to Idaho, New Mexico, Arizona and Nevada. ksacta Thickets and rocky ground[43]. Borders of streams, 900 - 2,500 metres in Texas[227].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. There is some doubt over the correct name for this species. [200] suggests that it is a synonym for *C. douglasii* but also records it as a distinct species[K]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.37 *Crataegus jackii*

4 tin 8 cyuc 2 kcaak 0 tcan 2 dwat 3 m hkuc
hmimhtut cyuti hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwlwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus jackii* is a deciduous Shrub growing to 3 m (9ft 10in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower in May. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[105, 177, 235]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is about 12mm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use.

only be consumed in very small quantities. Apple seeds do not normally contain very high quantities of hydrogen cyanide but, even so, should not be consumed in very large quantities. In small quantities, hydrogen cyanide has been shown to stimulate respiration and improve digestion, it is also claimed to be of benefit in the treatment of cancer. In excess, however, it can cause respiratory failure and even death.

kcaakta Edible Parts: FruitEdible Uses: PectinFruit - raw or cooked[159, 161]. Fairly large, it is up to 5cm in diameter[200]. Harsh and acid[11], it is mainly used for jellies[183] but can be eaten raw when it is fully ripe[101]. The fruits can be buried in the ground overwinter and will have lost much of their acidity by the spring[183]. The fruit can also be dried and stored for later use[257]. Rich in pectin, so it can be added to pectin-low fruits when making jams or jellies[226]. Pectin is also said to protect the body against radiation[201].

tcanta Pectin Rootstock WoodThe plant can be used as a rootstock for cultivated apples, conferring a greater hardness[226]. Wood - heavy, close-grained, not strong[46, 82, 229]. It weighs 43lb per cubic foot[235]. Used for making levers, the handles of tools, small domestic items and fuel[46, 82, 229].

dwatta Abortifacient Astringent Birthing aid Emmenagogue Lithontripic OphthalmicAn infusion of the bark has been used to ease a difficult birth and also in the treatment of gallstones, piles and as a wash for sore mouths[257]. A cold infusion of the bark has been used as a wash for black eyes, sore eyes and snow blindness[257]. A decoction of the root has been used to treat suppressed menses and so can cause an abortion, especially early in the pregnancy[257].

nrakta Eastern N. America. New York to South Carolina, west to Kansas. ksacta Bottoms, wooded slopes, thickets and clearings[43] in most soil types and moisture levels[159].

propta Seed - best sown as soon as it is ripe in the autumn in a cold frame. It usually germinates in late winter. Stored seed requires stratification for 3 months at 1°C and should be sown in a cold frame as soon as it is received[200]. It might not germinate for 12 months or more. Prick out the seedlings into individual pots as soon as they are large enough to handle. If given a rich compost they usually grow away quickly and can be large enough to plant out in late summer, though consider giving them some protection from the cold in their first winter. Otherwise, keep them in pots in a cold frame and plant them out in late spring of the following year. Cuttings of mature wood, November in a frame[11].

nwatta An easily grown plant, it succeeds in most fertile soils, preferring a moisture retentive well-drained loamy soil[1, 200]. Grows well in heavy clay soils. Prefers a sunny position but succeeds in partial shade, though it fruits less well in such a situation[200]. A slow-growing and short-lived tree in the wild[229]. The fruit is a good wildlife food source, especially for birds[200]. The flowers are sweetly scented[245]. This species is closely related to *M. ionensis*[1]. Plants in this genus are notably susceptible to honey fungus[200].

4.2.16 *Crataegus rotundifolia*

4 tin 8 cyuc 3 kcaak 0 tcan 2 dwat 6 m hkuc
hmimhtut cyuti hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwlwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus rotundifolia* is a deciduous Tree growing to 6 m (19ft) by 4 m (13ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in June, and the seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcaakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. A fair size fruit, to 15mm in diameter, with a reasonable sweet meaty taste, though when not fully ripe there is a distinct bitterness[K]. It makes an acceptable dessert fruit[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. There are up to five fairly

large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America. ksacta Not known

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A tree bearing this name at Kew in September 1993 was fruiting abundantly[K]. There is some confusion over the correct name of this species. *C. rotundifolia* Moench. is said to be a synonym for *C. chryscarpa* poenicea, *C. rotundifolia* Borckh. is a synonym of *C. chryscarpa*. There is also a *C. rotundifolia* Lam, which is a synonym of *Amelanchier rotundifolia*[200].

4.2.17 *Crataegus laevigata*

Midland Hawthorn, Smooth hawthorn, English Hawthorn
4 tin 8 cyuc 3 kcaak 5 tcan 5 dwat 6 m hkuc
pcimhmimhtut cyuti hgattcuphja7n hwicli hrasjlutnykfrip hpa7tli flatprikwlwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus laevigata* is a deciduous Shrub growing to 6 m (19ft) by 6 m (19ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from April to May, and the seeds ripen from September to November. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow

in full shade (deep woodland) semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: Fruit Leaves**Edible Uses:** Coffee TeaFruit - raw or cooked[2, 9, 12, 183]. A dry and mealy texture, they are not very appetizing[K]. The fruit can be used for jams and preserves[9]. The fruit pulp can be dried, ground into a meal and mixed with flour in making bread etc[46, 183]. The fruit is about 1cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K]. Young leaves and young shoots - raw[5, 177]. A tasty nibble, they are nice in a salad[K]. Young leaves are a tea substitute[21, 46, 177, 183]. The roasted seed is a coffee substitute[12, 21, 183].

tcanta Charcoal Fuel Hedge Hedge Rootstock WoodA good hedge plant, it is very tolerant of neglect and is able to regenerate if cut back severely, it makes a good thorny stock-proof barrier[186] and resists very strong winds. It can be used in layered hedges[11, 29]. The plant is often used as a rootstock for several species of garden fruit such as the medlar (*Mespilus germanica*) and the pear (*Pyrus communis sativa*) [4]. Wood - very hard and tough but difficult to work[7, 46, 61]. It has a fine grain and takes a beautiful polish but is seldom large enough to be of great value[4]. It is used for tool handles and making small wooden articles etc[4, 7, 46, 61]. The wood is valued in turning and makes an excellent fuel, giving out a lot of heat, more so even than oak wood[4]. Charcoal made from the wood is said to be able to melt pig iron without the aid of a blast[4].

dwatta Antiarrhythmic Antispasmodic Astringent Cardiotonic Diuretic Hypotensive Malaria Sedative Tonic VasodilatorHawthorn is an extremely valuable medicinal herb. It is used mainly for treating disorders of the heart and circulation system, especially angina[254]. Western herbalists consider it a 'food for the heart', it increases the blood flow to the heart muscles and restores normal heart beat[254]. This effect is brought about by the presence of bioflavonoids in the fruit, these bioflavonoids are also strongly antioxidant, helping to prevent or reduce degeneration of the blood vessels[254]. The fruit is antispasmodic, cardiac, diuretic, sedative, tonic and vasodilator[4, 9, 21, 46, 165]. Both the fruits and flowers of hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222], they are also used to treat a heart muscle weakened by age, for inflammation of the heart muscle, arteriosclerosis and for nervous heart problems[21]. Prolonged use is necessary for the treatment to be efficacious[222]. It is normally used either as a tea or a tincture[222]. Hawthorn is combined with ginkgo (*Ginkgo biloba*) to enhance poor memory, working by improving the blood supply to the brain[254]. The bark is astringent and has been used in the treatment of malaria and other fevers[7]. The roots are said to stimulate the arteries of the heart[218].

nrakta Europe, including Britain, from Sweden to Spain, eastwards to Poland. ksacta Woods, hedges, thickets etc on clays and heavy loams, especially in E. Britain[17, 186]. Where found in hedges it is often as a relic of ancient woodland[186].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be

best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta Landscape Uses:Border, Espalier, Pollard, Specimen, Street tree. A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200].. A true woodland species, it grows well in quite dense shade[17, 186]. A very hardy plant, tolerating temperatures down to at least -18°c[202]. Hybridizes freely with other members of this genus[200]. Closely allied to *C. monogyna*, it often hybridizes with that species in the wild when growing in its proximity[186]. There are many named forms selected for their ornamental value[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. An important food plant for the larvae of many lepidoptera species[30]. Special Features:Not North American native. Blooms are very showy.

4.2.18 Crataegus altaica

Altai Mountain Thorn

4 tlin 8 cyuc 3 kcak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikwis hpa7tmwatli xre7t hjonli

drutta Crataegus altaica is a deciduous Tree growing to 6 m (19ft) by 4 m (13ft). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Fruit - raw or cooked. About 8mm in diameter, the fruit is yellow with a fairly dry mealy texture and a pleasantly sweet flesh[K]. The fruit can also be used in making pies, preserves, etc, and can also be dried for later use. The fruit ripens in August, making it one of the earliest ripening hawthorns[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Although no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta W. Asia - Altai Mountains. ksacta Not known **propta** Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up

another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them out in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.34 Crataegus jonesiae

Miss jones' hawthorn

4 tlin 8 cyuc 2 kcak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikwissip hpa7tmwatli xre7t hjonli

drutta Crataegus jonesiae is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. The seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: Fruit**Edible Uses:** Fruit - raw or cooked. Sweet and mealy in one report[82], juicy according to others[11, 200]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is about 1.5cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta North-eastern N. America - Maine. ksacta Rocky shores of sounds and bays[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold

stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Closely related to *C. chrysoarpa* and considered to be no more than a form of that species by some botanists[229]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Hybridizes freely with other members of this genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.35 Crataegus canbyi

Cockspur hawthorn, Dwarf Hawthorn, Cockspur Hawthorn
4 tlin 7 cyuc 2 kcak 0 tcan 2 dwat 5 m hkuc hmimhtut cyutli hgatyuctcuphja7n hwicli hrasjlutnyikfrip hpa7tli flatprikwissip hpa7tmwatli xre7t hjonli

drutta Crataegus canbyi is a deciduous Shrub growing to 5 m (16ft) by 6 m (19ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in June, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: Fruit**Edible Uses:** Fruit - raw or cooked[82]. The fruit is about 12mm in diameter and is borne in small clusters[200]. It has a thick, bright red, juicy flesh[82]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - fine-grained, tough, hard and heavy. Used for tool handles etc[82].

dwatta Cardiac HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Pennsylvania and Delaware. ksacta Hedges and thickets[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year.

high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta South-eastern N. America - Virginia to Georgia, Illinois and Kansas. ksacta Thickets, open woods[43] and banks of streams in rich soils[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta Landscape Uses:Espalier, Firewood, Pollard, Specimen, Street tree. A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A very ornamental plant, there are some named varieties[1, 11, 200]. Special Features: North American native, Attractive flowers or blooms.

4.2.32 Malus florentina

Hawthorn-leaf crab apple

4 tlin 8 cyuc 2 kcaak 0 tcan 0 dwat 8 m hkuc hmimhtut cyutli tucphja7n hwicli hrasjltunykfrp hpa7tli flatprikliwslip hpa7tmwatli xre7t hjonli

drutta Malus florentina is a deciduous Tree growing to 8 m (26ft) by 6 m (19ft). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower in June, and the seeds ripen from October to November. The species is hermaphrodite (has both male and female organs) and is pollinated by Insects. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils, prefers well-drained soil and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil.

pcista All members of this genus contain the toxin hydrogen cyanide in their seeds and possibly also in their leaves, but not in their fruits. Hydrogen cyanide is the substance that gives almonds their characteristic taste but it should only be consumed in very small quantities. Apple seeds do not normally contain very high quantities of hydrogen cyanide but, even so, should not be consumed in very large quantities. In small quantities, hydrogen cyanide has been shown to stimulate respiration and improve digestion, it is also claimed to be of benefit in the treatment of cancer. In excess, however, it can cause respiratory failure and even death.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[K]. The oval fruit is about 15mm long and 8mm wide, it ripens in mid autumn and usually blets on the tree. When bletted, it has a mealy texture with a soft acid flesh, we found it refreshing in small quantities[K].

nrakta Europe - Northern Italy. ksacta Not known

propta Seed - best sown as soon as it is ripe in the autumn in a cold frame. It usually germinates in late winter. Stored seed requires stratification for 3 months at 1°c and should be sown in a cold frame as soon as it is received[200]. It might not germinate for 12 months or more. Prick out the seedlings into individual pots as soon as they are large enough to handle. If given a rich compost they usually grow away quickly and can be large enough to plant out in late summer, though consider giving them some protection from the cold in their first winter. Otherwise, keep them in pots in a cold frame and plant them out in late spring of the following year. Cuttings of mature wood, November in a frame[11].

nwatta An easily grown plant, it succeeds in most fertile soils, preferring a moisture retentive well-drained loamy soil[1, 200]. Grows well in heavy clay soils. Prefers a sunny position but succeeds in partial shade, though it fruits less well in such a situation[200]. This species is a very good pollinator for the cultivated apples[200]. The fruit is a good wildlife food source, especially for birds[200]. A very ornamental tree, especially when flowering in the spring, it makes a good specimen tree in a lawn[11]. Some botanists believe this species to be a bigeneric hybrid between *M. sylvestris* and *Sorbus torminalis*. They have assigned the name *Malosorbus florentina* to the plant, but this is not generally accepted[11]. Plants in this genus are notably susceptible to honey fungus[200].

4.2.33 Crataegus sanguinea

4 tlin 8 cyuc 2 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjltunykfrp hpa7tli flatprikliwslip hpa7tmwatli xre7t hjonli

drutta Crataegus sanguinea is a deciduous Shrub growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. It is in flower from May to June, and the seeds ripen from August to September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[2, 105, 177]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. It is about 1cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Hedge HedgeVery tolerant of trimming, it can even be cut right back into the old wood and will soon resprout. It makes a good hedge[74]. Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta E. Asia - northern China, Mongolia, Siberia. ksacta Open forests, forest edges, river banks etc[74].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°c and then cold stratified for another 3 months at 4°c[164]. It may still take

the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. This species is closely related to *C. wattiana*[11]. Hawthorns in general hybridize freely with other members of the genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.19 Crataegus flabellata

Fanleaf hawthorn

4 tlin 8 cyuc 3 kcaak 3 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjltunykfrp hpa7tli flatprikliwslip hpa7tmwatli xre7t hjonli

drutta Crataegus flabellata is a deciduous Shrub growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 4 and is not frost tender. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. A sweet but very soft and mealy flesh, it makes an acceptable dessert fruit[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is up to 15mm long and 8mm wide, it is borne in small clusters[82, 229]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Quebec to South Carolina, west to Louisiana. ksacta Thickets and open woods[204]. Dry open places, borders of woods and the margins of high banks of streams[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it

should be warm stratified for 3 months at 15°c and then cold stratified for another 3 months at 4°c[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. A tree at Kew in September 1993 was fruiting heavily[K].

4.2.20 Crataegus pruinosa

Frosted Hawthorn, Waxyfruit hawthorn

4 tlin 8 cyuc 3 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjltunykfrp hpa7tli flatprikliwslip hpa7tmwatli xre7t hjonli

drutta Crataegus pruinosa is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[105, 161, 177]. The flesh is thick[82]. A sweet yellow flesh[200]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use[257]. The fruit is about 1cm in diameter[200]. The fruit is up to 16mm in diameter with a thin flesh[229]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta North-eastern N. America - Newfoundland to North Carolina, west to Wisconsin and Oklahoma. ksacta

Thickets and rocky ground[43] in open woods[235]. Slopes of low hills, often in limestone soils[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.21 Crataegus monogyna

Hawthorn, Oneseed hawthorn

4 tin 8 cyuc 3 kcaak 3 tcan 5 dwat m hkuc hmimhtut cyutli hgattcuphja7n hwicli hyhrasj[lutnykifrip hpa7tli fruslatprikwvisslip hpa7tmwatli xre7t hjonli

drutta Crataegus monogyna is a deciduous Shrub growing to 6 m (19ft) by 6 m (19ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower from May to June, and the seeds ripen from September to November. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. It is noted for attracting wildlife. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very acid and very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: Flowers Fruit LeavesEdible Uses: Coffee TeaFruit - raw or cooked[2, 12]. Not very appetizing raw[9, K], it is normally used for making jams and preserves[9, 183]. The fruit can be dried, ground, mixed with flour and used for making bread etc[46]. The fruit is about 1cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K]. Young shoots - raw[5, 177]. A pleasant nutty flavour[144], they are a good addition to the salad bowl[183]. A tea is made from the dried leaves[21, 46, 177, 183], it is a china tea substitute. The roasted seeds are a coffee substitute[12, 21, 46, 177]. The flowers are used in syrups and sweet puddings[183].

tcanta Fuel Hedge Hedge WoodA good hedge plant, it is very tolerant of being cut and of neglect and is able to regenerate if cut back severely, it makes a good thorny stock-proof

barrier[186] and resists very strong winds. It is often used in layered hedges[11, 29]. The cultivar 'Stricta' has made a very good hedge 3.5 metres tall in an exposed maritime position at Rosewarne in N. Cornwall[K]. Wood - very hard and tough, difficult to work. Used for tool handles etc. Valued in turning[7, 46, 61]. A good fuel, giving out a lot of heat[4].

dwatta Antispasmodic Astringent Cardiotonic Diuretic Hypotensive Malaria Sedative Tonic VasodilatorHawthorn is an extremely valuable medicinal herb. It is used mainly for treating disorders of the heart and circulation system, especially angina[254]. Western herbalists consider it a 'food for the heart', it increases the blood flow to the heart muscles and restores normal heart beat[254]. This effect is brought about by the presence of bioflavonoids in the fruit, these bioflavonoids are also strongly antioxidant, helping to prevent or reduce degeneration of the blood vessels[254]. The fruit is antispasmodic, cardiac, diuretic, sedative, tonic and vasodilator[4, 9, 21, 46, 165]. Both the fruits and flowers of hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222], they are also used to treat a heart muscle weakened by age, for inflammation of the heart muscle, arteriosclerosis and for nervous heart problems[21]. Prolonged use is necessary for the treatment to be efficacious[222]. It is normally used either as a tea or a tincture[222]. Hawthorn is combined with ginkgo (Ginkgo biloba) to enhance poor memory, working by improving the blood supply to the brain[254]. The bark is astringent and has been used in the treatment of malaria and other fevers[7]. The roots are said to stimulate the arteries of the heart[218].

nrakta Europe, including Britain, absent from Iceland, south and west the the Mediterranean and Afghanistan ksacta Woods, hedges, thickets etc, on most soils except wet peat and poor acid sands[9, 17].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Succeeds in all but the very poorest acid soils[186]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. A very hardy plant, tolerating temperatures down to at least -18°C[202]. Hybridizes freely with other members of this genus and with C. laevigata in the wild[186, 200]. There are many named forms selected for their ornamental value[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being trans-

them on in the greenhouse or cold frame for their first winter. They can be planted out into their permanent positions in early summer of the following year, or be placed in an outdoor nursery bed for a year or so to increase in size. They might need protection from spring frosts. Cuttings of semi-ripe terminal shoots, 5 - 8cm long, August in a frame. Protect from frost. Forms roots in the spring[78]. Cuttings of mature terminal shoots, 5 - 10cm long, September/October in a cold frame. Takes 12 months[78]. Cuttings of soft to semi-ripe wood, early summer in a frame. Slow but sure.

nwatta Landscape Uses:Screen, Specimen. Likes abundant moisture at the roots, if grown in drier areas it must be given a deep moist soil[11]. Tolerates poor peaty soils[200]. Succeeds in wet cold and shallow soils but is not very wind-firm in shallow soils[1]. Prefers a pH between 4 to 6[200]. Another report says that it grows wild on very alkaline soils so it might be suitable for cultivation on alkaline soils. Dislikes shade[200]. Intolerant of atmospheric pollution[11]. Resists wind exposure to some degree[200]. In some upland areas, especially over granitic or other base-poor soils, growth rate and health have been seriously affected by aluminium poisoning induced by acid rain[200]. Plants are strongly outbreeding, self-fertilized seed usually grows poorly[200]. They hybridize freely with other members of this genus[200]. Fairly fast growing in Britain, able to average 30cm a year for at least the first 50 years[185]. Another report says that it is not a good grower in Britain[1]. Trees should be planted into their permanent positions when they are quite small, between 30 and 90cm. Larger trees will check badly and hardly put on any growth for several years. This also badly affects root development and wind resistance[200]. Special Features: Inconspicuous flowers or blooms.

4.2.30 Crataegus crus

Cockspur Thorn, Cockspur hawthorn, Dwarf Hawthorn 4 tin 7 cyuc 2 kcaak 0 tcan 2 dwat 10 m hkuc hmimhtut cyutli hgatyuctcuphja7n hwicli hrasj[lutnykifrip hpa7tli flatprikwvisslip hpa7tmwatli xre7t hjonli

drutta Crataegus crus-galli is a deciduous Shrub growing to 10 m (32ft) by 10 m (32ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in June, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[62, 101, 105]. A dry flesh, it is used in jellies[43]. The fruit is about 1cm in diameter and is borne in small clusters[200]. It often persists on the tree until spring[200]. This suggests that it does not make very good eating[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Hedge Hedge WoodVery amenable to trimming, the plant can even be cut right back into very old wood and will resprout freely. It is often used as a hedge[82]. Wood - fine-grained, hard and heavy. Used for tool handles etc[46, 61, 227].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Quebec to Georgia, west to Louisiana. Locally naturalized in Europe[50]. ksacta Thickets and open ground, especially in dry or rocky places[43]. Usually found on the slopes of low hills in rich soils[82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate

in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta Landscape Uses:Border, Espalier, Pollard, Screen, Specimen, Street tree. A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. A very hardy plant, tolerating temperatures down to about -18°C[202]. A very ornamental plant[1, 11]. There are many named forms, selected for their ornamental value[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. This plant is often confused in cultivation with C. prunifolia. 'Splendens'[11]. Special Features:Attracts birds, North American native, Blooms are very showy.

4.2.31 Crataegus phaenopyrum

Washington Thorn, Washington Hawthorn

4 tin 8 cyuc 2 kcaak 0 tcan 2 dwat 10 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasj[lutnykifrip hpa7tli flatprikwvisslip hpa7tmwatli xre7t hjonli

drutta Crataegus phaenopyrum is a deciduous Tree growing to 10 m (32ft) by 10 m (32ft) at a medium rate. See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in July, and the seeds ripen in October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[62]. A dry flesh[43]. Very small, about 5mm in diameter, it often hangs on the tree all winter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with

in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.28 *Crataegus maximowiczii*

4 tlin 8 cyuc 2 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjnutnyikfrip hpa7tli flatprikkwis hpa7tmwatli xre7t hjonli

drutta *Crataegus maximowiczii* is a deciduous Tree growing to 7 m (23ft). See above for USDA hardness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen from September to October. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[177]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. It is about 1cm in diameter[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta E. Asia - Siberia in N. Japan. ksacta Riverain deciduous forest strips, crests, meadows of flooded valleys, forest edges and dry mountain slopes[74].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year.

Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. This species is closely related to *C. sanguinea*[200].

4.2.29 *Picea glehnii*

Sakhalin Spruce

4 tlin 7 cyuc 2 kcaak 0 tcan 0 dwat 30 m hkuc htut cyutli tcuphja7n hwicli hyishrasjnutnyikfrip hpa7tli frusflatprikslip hpa7tmwatli hjonli

drutta *Picea glehnii* is an evergreen Tree growing to 30 m (98ft 5in) at a fast rate. See above for USDA hardness. It is hardy to UK zone 4. It is in leaf all year, and the seeds ripen from September to October. The species is monoecious (individual flowers are either male or female, but both sexes can be found on the same plant) and is pollinated by Wind. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay and nutritionally poor soils. Suitable pH: mildly acid and neutral soils and can grow in very acid and very alkaline soils. It cannot grow in the shade. It prefers moist or wet soil. The plant can tolerate strong winds but not maritime exposure. It cannot tolerate atmospheric pollution.

kcakta Edible Parts: Flowers Inner bark SeedEdible Uses: TeaYoung male catkins - raw or cooked. Used as a flavouring[172]. Immature female cones - cooked. The central portion, when roasted, is sweet and syrupy[172]. Inner bark - dried, ground into a powder and then used as a thickener in soups etc or added to cereals when making bread[172]. An emergency food, it is only used when all else fails. Seed - raw. Too small and fiddly to be worthwhile unless you are desperate[172]. A refreshing tea, rich in vitamin C, can be made from the young shoot tips[172].

tcanta WoodWood - beautifully grained. Used for pianos, violins, interiors of buildings etc[46, 61]. It is also valued for its use in the pulp industry to make paper[171].

nrakta E. Asia - N. Japan. ksacta Bogs in N. and C. Japan[58]. It is also found on very alkaline serpentine rocks.

propta Seed - stratification will probably improve germination so sow fresh seed in the autumn in a cold frame if possible[80]. Sow stored seed as early in the year as possible in a cold frame[78]. A position in light shade is probably best[78]. Seed should not be allowed to dry out and should be stored in a cool place[80]. Prick out the seedlings into individual pots when they are large enough to handle and grow

planted[11]. In heavier shade they quickly become drawn and leggy, eventually dying[186]. An important food plant for the caterpillars of many lepidoptera species[30], there are 149 insect species associated with this tree[24]. Plants are susceptible to fireblight[200]. In garden design, as well as the above-ground architecture of a plant, root structure considerations help in choosing plants that work together for their optimal soil requirements including nutrients and water. The root pattern is branching: a heart root, dividing from the crown into several primary roots going down and out [2-1].

4.2.22 *Crataegus chrysoarpa*

Fireberry Hawthorn, Red haw, Piper's hawthorn, 4 tlin 8 cyuc 3 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgattcuphja7n hwicli hrasjnutnyikfrip hpa7tli flatprikkwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus chrysoarpa* is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[161, 177]. Used mainly as a famine food[257]. A very pleasant flavour when ripe, with the added bonus of ripening in late summer before most other members of the genus[K]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. It is about 1cm in diameter and borne in small clusters[200]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K]. A tea can be made from the twigs[257]. (This probably means the young shoots with leaves[K].)

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic Hypotensive LaxativeA decoction of the dried berries has been used as a mild laxative[257]. A compound decoction of the root has been used in the treatment of diarrhoea[257]. Although no other specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta North-eastern N. America - Newfoundland to Pennsylvania, west to the Rocky Mountains. ksacta Thickets and rocky ground along streams[43].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but

undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. A ten year old tree was seen at Kew Gardens in 2002. It was about 2.5 metres tall and was bearing a very good crop of fruit[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Hybridizes freely with other members of this genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.23 *Crataegus coccinoides*

Kansas Hawthorn

4 tlin 8 cyuc 3 kcaak 0 tcan 2 dwat 6 m hkuc hmimhtut cyutli hgatkuytucphja7n hwicli hrasjnutnyikfrip hpa7tli flatprikkwisslip hpa7tmwatli xre7t hjonli

drutta *Crataegus coccinoides* is a deciduous Tree growing to 6 m (19ft 8in). See above for USDA hardness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers dry moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[2, 177]. Firm and sub-acid[82]. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. The fruit is borne in small clusters and is up to 17mm in diameter[229]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Central N. America - Illinois and Missouri to Kansas, Oklahoma and Arkansas. ksacta Dry thickets and calcareous hills[43].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their

final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Hybridizes freely with other members of this genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.24 Crataegus anomala

Arnold hawthorn

4 tlin 8 cyuc 3 kcak 3 tcan 2 dwat 5 m hkuc
hminhtut cyutli hgattcuphja7n hwicli hrasljutnyikfrip hpa7tli
flatprikwiwslip hpa7tmwatli xre7t hjonli

drutta Crataegus anomala is a deciduous Shrub growing to 5 m (16ft 5in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[105, 177, 235]. Juicy[43]. A very nice flavour; it makes a very good dessert fruit[K]. The fruit can also be used in making pies, preserves, etc, and can be dried for later use. The fruit is up to 20mm in diameter with a thick flesh[235]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Quebec to New York. ksacta Rocky banks and open woods on low limestone ridges[43, 82].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to

pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. This species is growing in a number of botanical gardens in Britain, where it is fruiting very well[K]. Some botanists do not recognise it as a distinct species and place it as part of *C. pedicellata*[82]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11].

4.2.25 Crataegus columbiana

Columbian Hawthorn

4 tlin 8 cyuc 3 kcak 0 tcan 2 dwat 5 m hkuc
hminhtut cyutli hgattcuphja7n hwicli hrasljutnyikfrip hpa7tli
flatprikwiwslip hpa7tmwatli xre7t hjonli

drutta Crataegus columbiana is a deciduous Tree growing to 5 m (16ft 5in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked[161, 166]. A very pleasant flavour, it is slightly mealy but juicy and can be eaten in quantity as a dessert fruit[K]. It is also used in making pies, preserves etc and can be dried for later use[161, 177]. The fruit is up to 11mm in diameter and is borne in small clusters[K]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Western N. America - British Columbia to California, east to Idaho and Oregon. ksacta Meadows and near streams in California[71].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months at 4°C[164]. It may still take

another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. This species is closely related to *C. douglasii*[71].

4.2.26 Crataegus x grignonensis

4 tlin 8 cyuc 3 kcak 3 tcan 2 dwat 4 m hkuc
hminhtut cyutli hgattcuphja7n hwicli hrasljutnyikfrip hpa7tli
flatprikwiwslip hpa7tmwatli xre7t hjonli

drutta Crataegus x grignonensis is a deciduous Shrub growing to 4 m (13ft 1in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw or cooked. The fruit can be used in making pies, preserves, etc, and can also be dried for later use. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta A hybrid species of garden origin, *C. crus-galli* x *C. pubescens*. ksacta Not known in the wild.

propta Seed - this is a hybrid species and it will not breed true from seed. If this is no problem, then the seed is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate in the spring, though most will probably take another year. Stored seed can be very slow and erratic to germinate, it should be warm stratified for 3 months at 15°C and then cold stratified for another 3 months

at 4°C[164]. It may still take another 18 months to germinate[78]. Scarifying the seed before stratifying it might reduce this time[80]. Fermenting the seed for a few days in its own pulp may also speed up the germination process[K]. Another possibility is to harvest the seed 'green' (as soon as the embryo has fully developed but before the seedcoat hardens) and sow it immediately in a cold frame. If timed well, it can germinate in the spring[80]. If you are only growing small quantities of plants, it is best to pot up the seedlings as soon as they are large enough to handle and grow them on in individual pots for their first year, planting them out in late spring into nursery beds or their final positions. When growing larger quantities, it might be best to sow them directly outdoors in a seedbed, but with protection from mice and other seed-eating creatures. Grow them on in the seedbed until large enough to plant out, but undercut the roots if they are to be left undisturbed for more than two years.

nwatta A very easily grown plant, it prefers a well-drained moisture retentive loamy soil but is not at all fussy[11, 200]. Once established, it succeeds in excessively moist soils and also tolerates drought[200]. It grows well on a chalk soil and also in heavy clay soils[200]. A position in full sun is best when plants are being grown for their fruit, they also succeed in semi-shade though fruit yields and quality will be lower in such a position[11, 200]. Most members of this genus succeed in exposed positions, they also tolerate atmospheric pollution[200]. Hybridizes freely with other members of this genus[200]. Seedlings should not be left in a seedbed for more than 2 years without being transplanted[11]. Seedling trees take from 5 - 8 years before they start bearing fruit, though grafted trees will often flower heavily in their third year[K]. The flowers have a foetid smell somewhat like decaying fish. This attracts midges which are the main means of fertilization. When freshly open, the flowers have more pleasant scent with balsamic undertones[245].

4.2.27 Crataegus intricata

Copenhagen hawthorn

4 tlin 8 cyuc 3 kcak 3 tcan 2 dwat 3 m hkuc
hminhtut cyutli hgattcuphja7n hwicli hrasljutnyikfrip hpa7tli
flatprikwiwslip hpa7tmwatli xre7t hjonli

drutta Crataegus intricata is a deciduous Shrub growing to 3 m (9ft 10in). See above for USDA hardiness. It is hardy to UK zone 5 and is not frost tender. It is in flower in May, and the seeds ripen in September. The species is hermaphrodite (has both male and female organs) and is pollinated by Midges. Suitable for: light (sandy), medium (loamy) and heavy (clay) soils and can grow in heavy clay soil. Suitable pH: mildly acid, neutral and basic (mildly alkaline) soils and can grow in very alkaline soils. It can grow in semi-shade (light woodland) or no shade. It prefers moist or wet soil and can tolerate drought. The plant can tolerate strong winds but not maritime exposure. It can tolerate atmospheric pollution.

kcakta Edible Parts: FruitEdible Uses: Fruit - raw, cooked or dried for later use[2, 105, 161]. Hard dry flesh[43]. Fruit we have eaten from this tree is mealy and fairly soft, with a pleasant sweet flavour[K]. It ripens in early September in southern Britain[K]. The fruit is up to 14mm in diameter and is borne in small clusters[229]. The skin is slightly hairy[235]. There are up to five fairly large seeds in the centre of the fruit, these often stick together and so the effect is of eating a cherry-like fruit with a single seed[K].

tcanta Wood - heavy, hard, tough, close-grained. Useful for making tool handles, mallets and other small items[82].

dwatta Cardiotonic HypotensiveAlthough no specific mention has been seen for this species, the fruits and flowers of many hawthorns are well-known in herbal folk medicine as a heart tonic and modern research has borne out this use. The fruits and flowers have a hypotensive effect as well as acting as a direct and mild heart tonic[222]. They are especially indicated in the treatment of weak heart combined with high blood pressure[222]. Prolonged use is necessary for it to be efficacious[222]. It is normally used either as a tea or a tincture[222].

nrakta Eastern N. America - Maine to Georgia, west to Michigan and Oklahoma. ksacta Thickets and open rocky woods[43, 235].

propta Seed - this is best sown as soon as it is ripe in the autumn in a cold frame, some of the seed will germinate